



A DOCPHOENIX

☐ TRNA _____
Transmittal New Application

☐ SPEC _____
Specification

☐ CLM _____
Claims

☐ ABST _____
Abstract

☐ DRW _____
Drawings

☐ OATH _____
Oath or Declaration

☐ ADS _____
Application Data Sheet

☐ A... _____
Amendment Including Elections

☐ A.PE _____
Preliminary Amendment

☐ REM _____
Applicant Remarks in Amendment

☐ IDS _____
IDS Including 1449


☐ 371P _____
PCT Papers in a 371P Application

☒ FOR 214 _____
Foreign Reference

☐ NPL _____
Non-Patent Literature

☐ FRPR _____
Foreign Priority Papers

☐ ARTIFACT _____
Artifact



☐ TRREISS _____
Transmittal New Reissue Application

☐ PROTRANS _____
Translation of Provisional in Nonprovisional

☐ BIB _____
Bib Data Sheet

☐ WCLM _____
Claim Worksheet

☐ WFEE _____
Fee Worksheet

☐ APPENDIX _____
Appendix

☐ COMPUTER _____
Computer Program Listing

☐ SPEC NO _____
Specification Not in English

☐ N417 _____
Copy of EFS Receipt Acknowledgement

☐ CRFL _____
Computer Readable Form Transfer Request Filed

☐ CRFS _____
Computer Readable Form Statement

☐ SEQLIST _____
Sequence Listing

☐ SIR. _____
SIR Request

☐ AF/D _____
Affidavit or Exhibit Received

☐ DIST _____
Terminal Disclaimer Filed

☐ PET. _____
Petition

☐ END JOB☐ DUPLEX

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 June 2002 (20.06.2002)

PCT

(10) International Publication Number
WO 02/47690 A1

- (51) International Patent Classification⁷: **A61K 31/50**, 31/53, 31/505
- (21) International Application Number: PCT/US01/47498
- (22) International Filing Date:
12 December 2001 (12.12.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/254,581 12 December 2000 (12.12.2000) US
- (71) Applicant: **CYTOVIA, INC.** [US/US]; 6650 Nancy Ridge Drive, San Diego, CA 92121 (US).
- (72) Inventors: **CAI, Sui, Xiong**; 3623 Berryfield Court, San Diego, CA 92130 (US). **DREWE, John, A.**; 7069 Murillo Lane, Carlsbad, CA 92009 (US). **NGUYEN, Bao**; 9494 Carroll Canyon Road #28, San Diego, CA 92126 (US). **REDDY, P, Sanjeeva**; 13293 Larkfield Court, San Diego, CA 92130 (US). **PERVIN, Azra**; 946 Calle Capistrano, San Marcos, CA 92069 (US).
- (74) Agents: **ESMOND, Robert, W.** et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/47690 A1

(54) Title: SUBSTITUTED 2-ARYL-4-ARYLAMINOPYRIMIDINES AND ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF APOPTOSIS AND THE USE THEREOF

(57) Abstract: The present invention is directed to substituted 2-aryl-4-arylamino-pyrimidine and analogs thereof, represented by the general Formula I: wherein A, Ar₁, Ar₂, and R₁ and R₂ are defined herein. The present invention also relates to the discovery that compounds having Formula I are activators of caspases and inducers of apoptosis. The compounds of this invention may be used to induce cell death in a variety of clinical conditions in which uncontrolled growth and spread of abnormal cells occurs.

SUBSTITUTED 2-ARYL-4-ARYLAMINOPYRIMIDINES AND
ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF
APOPTOSIS AND THE USE THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention is in the field of medicinal chemistry. In particular, the invention relates to substituted 2-aryl-4-arylamino-pyrimidines and analogs, and the discovery that these compounds are activators of caspases and inducers of apoptosis. The invention also relates to the use of these compounds as therapeutically effective anti-cancer agents.

Related Art

[0002] Organisms eliminate unwanted cells by a process variously known as regulated cell death, programmed cell death or apoptosis. Such cell death occurs as a normal aspect of animal development as well as in tissue homeostasis and aging (Glucksmann, A., *Biol. Rev. Cambridge Philos. Soc.* 26:59-86 (1951); Glucksmann, A., *Archives de Biologie* 76:419-437 (1965); Ellis, *et al.*, *Dev.* 112:591-603 (1991); Vaux, *et al.*, *Cell* 76:777-779 (1994)). Apoptosis regulates cell number, facilitates morphogenesis, removes harmful or otherwise abnormal cells and eliminates cells that have already performed their function. Additionally, apoptosis occurs in response to various physiological stresses, such as hypoxia or ischemia (PCT published application WO96/20721).

[0003] There are a number of morphological changes shared by cells experiencing regulated cell death, including plasma and nuclear membrane blebbing, cell shrinkage (condensation of nucleoplasm and cytoplasm), organelle relocalization and compaction, chromatin condensation and production of apoptotic bodies (membrane enclosed particles containing intracellular material) (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

[0004] Apoptosis is achieved through an endogenous mechanism of cellular suicide (Wyllie, A.H., in *Cell Death in Biology and Pathology*, Bowen and

- 2 -

Lockshin, eds., Chapman and Hall (1981), pp. 9-34). A cell activates its internally encoded suicide program as a result of either internal or external signals. The suicide program is executed through the activation of a carefully regulated genetic program (Wyllie, *et al.*, *Int. Rev. Cyt.* 68:251 (1980); Ellis, *et al.*, *Ann. Rev. Cell Bio.* 7:663 (1991)). Apoptotic cells and bodies are usually recognized and cleared by neighboring cells or macrophages before lysis. Because of this clearance mechanism, inflammation is not induced despite the clearance of great numbers of cells (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

[0005] It has been found that a group of proteases are a key element in apoptosis (see, e.g., Thornberry, *Chemistry and Biology* 5:R97-R103 (1998); Thornberry, *British Med. Bull.* 53:478-490 (1996)). Genetic studies in the nematode *Caenorhabditis elegans* revealed that apoptotic cell death involves at least 14 genes, two of which are the pro-apoptotic (death-promoting) *ced* (for *cell death abnormal*) genes, *ced-3* and *ced-4*. CED-3 is homologous to interleukin 1 beta-converting enzyme, a cysteine protease, which is now called caspase-1. When these data were ultimately applied to mammals, and upon further extensive investigation, it was found that the mammalian apoptosis system appears to involve a cascade of caspases, or a system that behaves like a cascade of caspases. At present, the caspase family of cysteine proteases comprises 14 different members, and more may be discovered in the future. All known caspases are synthesized as zymogens that require cleavage at an aspartyl residue prior to forming the active enzyme. Thus, caspases are capable of activating other caspases, in the manner of an amplifying cascade.

[0006] Apoptosis and caspases are thought to be crucial in the development of cancer (*Apoptosis and Cancer Chemotherapy*, Hickman and Dive, eds., Humana Press (1999)). There is mounting evidence that cancer cells, while containing caspases, lack parts of the molecular machinery that activates the caspase cascade. This makes the cancer cells lose their capacity to undergo cellular suicide and the cells become cancerous. In the case of the apoptosis process, control points are known to exist that represent points for intervention

- 3 -


leading to activation. These control points include the CED-9-BCL-like and CED-3-ICE-like gene family products, which are intrinsic proteins regulating the decision of a cell to survive or die and executing part of the cell death process itself, respectively (see, Schmitt, *et al.*, *Biochem. Cell. Biol.* 75:301-314 (1997)). BCL-like proteins include BCL-xL and BAX-alpha, which appear to function upstream of caspase activation. BCL-xL appears to prevent activation of the apoptotic protease cascade, whereas BAX-alpha accelerates activation of the apoptotic protease cascade.

[0007] It has been shown that chemotherapeutic (anti-cancer) drugs can trigger cancer cells to undergo suicide by activating the dormant caspase cascade. This may be a crucial aspect of the mode of action of most, if not all, known anticancer drugs (Los, *et al.*, *Blood* 90:3118-3129 (1997); Friesen, *et al.*, *Nat. Med.* 2:574 (1996)). The mechanism of action of current antineoplastic drugs frequently involves an attack at specific phases of the cell cycle. In brief, the cell cycle refers to the stages through which cells normally progress during their lifetimes. Normally, cells exist in a resting phase termed G₀. During multiplication, cells progress to a stage in which DNA synthesis occurs, termed S. Later, cell division, or mitosis occurs, in a phase called M. Antineoplastic drugs such as cytosine arabinoside, hydroxyurea, 6-mercaptopurine, and methotrexate are S phase specific, whereas antineoplastic drugs such as vincristine, vinblastine, and paclitaxel are M phase specific. Many slow growing tumors, for example colon cancers, exist primarily in the G₀ phase, whereas rapidly proliferating normal tissues, for example bone marrow, exist primarily in the S or M phase. Thus, a drug like 6-mercaptopurine can cause bone marrow toxicity while remaining ineffective for a slow growing tumor. Further aspects of the chemotherapy of neoplastic diseases are known to those skilled in the art (*See, e.g., Hardman, et al., eds., Goodman and Gilman's The Pharmacological Basis of Therapeutics*, Ninth Edition, McGraw-Hill, New York (1996), pp. 1225-1287). Thus, it is clear that the possibility exists for the activation of the caspase cascade, although the exact mechanisms for doing so are not clear at this point. It is equally

- 4 -

clear that insufficient activity of the caspase cascade and consequent apoptotic events are implicated in various types of cancer. The development of caspase cascade activators and inducers of apoptosis is a highly desirable goal in the development of therapeutically effective antineoplastic agents. Moreover, since autoimmune disease and certain degenerative diseases also involve the proliferation of abnormal cells, therapeutic treatment for these diseases could also involve the enhancement of the apoptotic process through the administration of appropriate caspase cascade activators and inducers of apoptosis.

[0008] Eur. Pat. Appl. EP 407899 discloses aminopyrimidine derivatives with activity as fungicide:



wherein,

R₁ is H, alkyl, alkoxyalkyl, alkylthioalkyl, cycloalkyl, alkenyl, alkynyl, cycloalkylalkyl, substituted aminoalkyl, phenyl, phenylalkyl, phenoxyalkyl, phenylmercaptoalkyl or phenoxyphenoxyalkyl, wherein the phenyl-portions are optionally substituted;

R₂, R₃, R₄ independently are H, alkyl or optionally substituted phenyl;

R₅ is H, alkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, alkoxy, alkylthio, alkoxyalkyl, R₇R₈N-, alkylthioalkyl, R₇R₈-alkyl, halogen, alkenyl, alkynyl, phenyl, phenoxy, phenylalkyl, phenoxyalkyl, phenylmercaptoalkyl, phenylmercapto, phenylalkoxy or phenylalkylthio, wherein the phenyl-portions are optionally substituted;

R₆ is H, alkyl, alkoxy, alkenyloxy, alkynyloxy, alkylthio, halogen or optionally substituted phenyl; or


R₅ and R₆ are taken together to form a polymethylene group;

- 5 -

R₇ and R₈ independently are H, alkyl, alkoxyalkyl, hydroxyalkyl, alkylthioalkyl, alkenyl, substituted aminoalkyl, alkynyl, cycloalkyl, cycloalkylalkyl, which in the cycloalkyl-portion is optionally substituted, formyl, phenyl or phenylalkyl, which in the phenyl-portion is optionally substituted; or


R₇ and R₈ are taken together with the nitrogen atom to which they are attached to form an optionally substituted 5- to 7-member, saturated or unsaturated, heterocycle with 1 or 3 heteroatoms, which are the same or different; and the acid addition salts which are functional as fungicides.

[0009] WO 0127089 patent application discloses pyrimidine derivatives for the treatment of diseases or medical conditions mediated by cytokines:



wherein,

m is 0, 1, 2 or 3 are each R¹ group, which may be the same or different, is selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, sulphamoyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or from a group of the formula:



wherein X¹ is a direct bond or is selected from O, S, SO, SO₂, N(R⁴), CO, CH(OR⁴), CON(R⁴), N(R⁴)CO, SO₂N(R⁴), N(R⁴), N(R⁴)SO₂, OC(R⁴)₂,

- 6 -

$SC(R^4)_2$ and $N(R^4)C(R^4)_2$, wherein each R^4 is hydrogen or (1-6C)alkyl, and Q^2 is aryl-(1-6C)alkyl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, or $(R^1)_m$ is (1-3C)alkylenedioxy, and wherein any aryl, heteroaryl or heterocyclyl group within a substituted on R^1 optionally bears 1, 2 or 3 substituents,

R_3 is hydrogen, halogeno or (1-6C)alkyl;

n is 0, 1 or 2 and each R^2 group, which may be the same or different is selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, (1-6C)alkoxycarbonyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di[(1-6C)alkyl]amino;

p is 0, 1, 2, 3, or 4; and

Q^1 is aryl or heteroaryl and Q^1 is optionally substituted with 1, 2, or 3 substituents, which may be the same or different, selected from hydroxy, halogeno, trifluoromethyl, cyano, mercapto, nitro, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C) alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(2-6C)alkyl-(2-6C)alkanoylamino, N-(2-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino and N-(1-6C)alkyl-(1-6C)alkanesulphonylamino, or with a (1-3C)alkylenedioxy group, or from a group of the formula:




wherein X^3 is a direct a bond or is selected from 0 and $N(R^8)$, wherein R^8 is hydrogen or (1-6C)alkyl, and Q^4 is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)-alkyl, heterocyclyl or heterocyclyl(1-6C)alkyl, and any Q^4 group optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, hydroxy, amino, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino.

[0010] WO 0027824 patent application discloses substituted pyrimidine compositions and methods of use. The compounds are said to have activity as

- 7 -


inhibitors of phospholipase A₂, and are useful in treating disorders mediated by phospholipase A₂:



wherein,

the symbol R₁ represents a C₁-C₆ alkyl, C₁-C₆ alkoxy or halogen atom. The symbol R₂ represents a phenyl group, substituted phenyl group, benzyl moiety, substituted benzyl moiety, C₃-C₇ cycloalkyl, or substituted C₃-C₇ cycloalkyl. The symbol R₃ represents a hydrogen or C₁-C₆ alkyl group. The symbol R₄ represents -H, -OH, -N₃ or -NHCOCH₃. The symbol R₅ represents H or alkyl, preferably H.

[0011] US patent 6,156,755 discloses the use of pyridine derivatives for the prevention of cancer:



wherein,

R¹ is hydrogen, halogen, cyano, nitro, trifluoromethyl, amino, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy, (C₆-C₁₂)-aryl, (C₁-C₆)-alkoxycarbonyl-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl-S-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl-SO-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl-SO₂-(C₁-C₆)-alkyl, dihydroxy-(C₁-C₆)-alkyl, aryl, heteroaryl, heteroaryl-(C₁-C₆)-alkyl, aryl-(C₁-C₆)-alkyl, (C₁-C₆)-alkoxycarbonylaryl, aryl-(C₁-C₆)-alkyloxy or heteroaryl-(C₁-C₆)-alkyloxy, heteroaryl is pyridyl, furyl, tetrahydrofuryl, thienyl, imidazolyl, pyrazolyl, triazolyl, thiazolyl, oxazolyl, benzothiazolyl;

- 8 -

R^2 , R^3 independently of one another are hydrogen, (C₁-C₆)-alkyl, (C₆-C₁₂)-aryl, (C₆-C₁₂)-arylalkyl having 1-4 alkyl carbon atoms, where aryl can be substituted by one to three substituents selected from the group consisting of chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, or

R^2 , R^3 , together with the nitrogen to which they are bonded, form the azetidino, pyrrolidino, piperidino, piperazino or morpholino group, where the heterocycles can be substituted by one or two substituents selected from the group consisting of chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -S-(C₁-C₆)-alkyl, -SO-(C₁-C₆)-alkyl, -SO₂-(C₁-C₆)-alkyl, sulfamoyl, N-(C₁-C₄)-alkylsulfamoyl, N,N-(C₁-C₄)-dialkylsulfamoyl, (C₁-C₆)-alkoxycarbonyl, N,N-(C₁-C₄)-dialkylcarbamoyl, N-(C₁-C₄)-alkylcarbamoyl, N-(C₆-C₁₂)-arylcarbamoyl, (C₆-C₁₂)-arylcarbamoyl substituted in the aryl radical by (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, halogen, NO₂, NH₂, CN or CF₃, (C₆-C₁₂)-arylcarbonyl substituted in the aryl radical by (C₁-C₄)-alkoxy, halogen, NO₂, NH₂, CN or CF₃, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, (C₆-C₁₂)-arylsulfonyl, (C₆-C₁₂)-arylsulfonyl substituted in the aryl radical by (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, halogen, NO₂, NH₂, CN or CF₃, heteroarylcarbonyl or heteroarylsulfonyl;

R^4 and R^5 independently of one another are hydrogen, halogen, cyano, nitro, trifluoromethyl, amino, (C₁-C₆)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₁-C₆)-alkoxy, (C₆-C₁₂)-aryl, naphthyl, furyl, where (C₆-C₁₂)-aryl, naphthyl and furyl can be substituted by one or two substituents selected from the group consisting of chlorine, bromine, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, -S-(C₁-C₆)-alkyl, -SO-(C₁-C₆)-alkyl, -SO₂-(C₁-C₆)-alkyl, hydroxyl; and their physiologically tolerable salts.


SUMMARY OF THE INVENTION

[0012] The present invention is related to the discovery that substituted 2-aryl-4-arylaminopyrimidines and analogs, as represented in Formula I, are activators of the caspase cascade and inducers of apoptosis. Thus, an aspect of

- 9 -

the present invention is directed to the use of compounds of Formula I as inducers of apoptosis.

[0013] Compounds of the present invention are represented by Formula I:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

Ar₁ and Ar₂ are independently and optionally substituted aryl or heteroaryl;

A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, an optionally substituted alkyl or cycloalkyl.

[0014] A second aspect of the present invention is to provide a method for treating, preventing or ameliorating neoplasia and cancer by administering a compound of Formula I to a mammal in need of such treatment.

[0015] Many of the compounds within the scope of the present invention are novel compounds. Therefore, a third aspect of the present invention is to provide novel compounds of Formula I, and to also provide for the use of these novel compounds for treating, preventing or ameliorating neoplasia and cancer.

[0016] A fourth aspect of the present invention is to provide a pharmaceutical composition useful for treating disorders responsive to the induction of apoptosis, containing an effective amount of a compound of Formula I in admixture with one or more pharmaceutically acceptable carriers or diluents.

- 10 -

[0017] A fifth aspect of the present invention is directed to methods for the preparation of novel compounds of Formula I.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES


[0018] Figs. 1A-B are graphs showing drug induced cell cycle arrest and apoptosis in T47D cells. Fig. 1A: control cells showing most of the cells in G1 phase of the cell cycle (M2). Fig. 1B: cells treated with 200 nM of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine for 48 h showing a reduction in the G1 population (M2), an increase in the G2/M population (M4) and the sub-diploid DNA population of cells (M1).

[0019] Fig. 2 is a graph showing inhibition of clonogenic survival of MX-1 and T47D cells treated for 48 h with different concentrations of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine. Fig. 2 shows increasing inhibition of clonogenicity with increasing drug concentration, with IC50 of about 100 nM and 300 nM for T47D and MX-1 cells, respectively.

DETAILED DESCRIPTION OF THE INVENTION

[0020] The present invention arises out of the discovery that substituted 2-aryl-4-arylaminopyrimidines and analogs, as represented in Formula I, are potent and highly efficacious activators of the caspase cascade and inducers of apoptosis. Therefore compounds of Formula I are useful for treating disorders responsive to induction of apoptosis.

[0021] Specifically, compounds of the present invention are represented by Formula I:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 11 -

Ar₁ and Ar₂ are independently and optionally substituted aryl or heteroaryl;


A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, an optionally substituted alkyl or cycloalkyl.

[0022] Preferred compounds of Formula I include compounds wherein Ar₁ or Ar₂ is optionally substituted phenyl, naphthyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolyl, isoquinolyl, thienyl, furyl, or pyrrolyl. Preferred compounds of Formula I also include compounds wherein R₃ is hydrogen. Preferred compounds of Formula I also include compounds wherein A is C-R₂. Especially preferred compounds of Formula I include compounds wherein Ar₁ is optionally substituted pyridinyl, pyrimidinyl and pyrazinyl. Especially preferred compounds of Formula I also include compounds wherein Ar₂ is optionally substituted phenyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl.

[0023] Preferred structures of Formula I are substituted 2-aryl-4-arylamino pyrimidines and analogs represented by Formula II:



or pharmaceutically acceptable salts or prodrugs thereof, wherein R₁-R₃, and Ar₂ are as defined in Formula I;

B is N or C-R₄;

D is N or C-R₅;

E is N or C-R₆;

F is N or C-R₇;


G is N or C-R₈; and

- 12 -

R₄-R₈ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; provided that not more than three of B, D, E, F and G are N.

[0024] Preferred compounds falling within the scope of Formula II include compounds wherein R₁ is an optionally substituted alkyl, haloalkyl or phenyl. Preferred compounds of Formula II also include compounds wherein Ar₂ is an optionally substituted phenyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl. Preferred compounds of Formula II also include compounds wherein B is nitrogen, D is C-R₅, E is C-R₆, F is C-R₇, and G is C-R₈. Preferred compounds of Formula II also include compounds wherein D is nitrogen, B is C-R₄, E is C-R₆, F is C-R₇, and G is C-R₈. Preferred compounds of Formula II also include compounds wherein E is nitrogen, D is C-R₅, B is C-R₄, F is C-R₇, and G is C-R₈. Preferred compounds of Formula II also include compounds wherein two of B, D, E, F and G are N.

[0025] Preferred structures of Formula I are substituted 2-aryl-4-arylamino-pyrimidines and analogs represented by Formula III:



or pharmaceutically acceptable salts or prodrugs thereof, wherein R₁-R₃, and B, D, E, F and G are as defined in Formula I and II;

R₉-R₁₃ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl,

arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol.

[0026] Preferred compounds falling within the scope of Formula III include compounds wherein R_1 is an optionally substituted alkyl, haloalkyl. Preferred compounds of Formula III also include compounds wherein R_2 is hydrogen. Preferred compounds of Formula III also include compounds wherein one of the B, D, E, F and G is nitrogen. Preferred compounds of Formula III also include compounds wherein two of the B, D, E, F and G is nitrogen. Preferred compounds of Formula III also include compounds wherein R_{10} or R_{12} are not hydrogen. Preferred compounds of Formula III also include compounds wherein R_9 and R_{12} are not hydrogen. Preferred compounds of Formula III also include compounds wherein R_{10} and R_{12} are not hydrogen. Preferred compounds of Formula III also include compounds wherein R_9 , R_{11} and R_{12} are not hydrogen.

[0027] Exemplary preferred compounds that may be employed in the method of the invention include, without limitation:

4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dichloroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4-Difluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methylanilino)-2-phenyl-6-methylpyrimidine;
4-(4-Methoxyanilino)-2-(2-hydroxyphenyl)-6-methylpyrimidine;
4-(4-(Trifluoromethoxy)anilino)-2-(pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 14 -

4-(3-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-(Trifluoromethoxy)anilino)-2-(2-pyridinyl)-6-t-butylpyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-t-butylpyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-chloropyrimidine;
4-(4-Methoxyanilino)-2-phenyl-6-chloropyrimidine;
4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(4-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3,4,5-Trimethoxyanilino)-2-(N-pyrrolyl)-1,3,5-triazine;
4-(3,5-Dichloroanilino)-2-(N-pyrrolyl)-1,3,5-triazine;
4-[2-(Trifluoromethyl)benzylamino]-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 15 -

4-(4-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
4-(2-Chloroanilino)-2-(N-pyrrolyl)-1,3,5-triazine;
4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 16 -

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

- 17 -

4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 18 -

4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine
hydrochloride;
4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine
hydrochloride;
4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-
(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride;
4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-
(trifluoromethyl)pyrimidine;
4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-
(trifluoromethyl)pyrimidine hydrochloride;

- 19 -

4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine; 4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

- 20 -

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide;

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 21 -

4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(Indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine;

4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(4-Fluoro-3-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-pyridinyl)pyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

- 22 -

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

- 23 -

4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;
 4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
 4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride;
 4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(t-butyl)pyrimidine;
 4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine;
 4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine;
 4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
 4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine; and
 4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine.

[0028] The present invention is also directed to novel compounds within the scope of Formulae I-III. Exemplary preferred compounds that may be employed in this invention include, without limitation:

4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(4-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(4-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
 4-(3-Benzyloxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Ethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Cyanoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Acetophenonanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 4-(3-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
 4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

- 24 -

4-[3-(Trifluoromethyl)anilino]-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Acetoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Nitroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-[3-(Trifluoromethoxy)anilino]-6-methyl-2-(2-pyridinyl)pyrimidine;
4-[3-(Methylthio)anilino]-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(3-Methoxy-phenoxy)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-5-methoxy-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
6-Morpholino-4-(3-methoxyanilino)-2-phenyl-pyrimidine;
6-Morpholino-4-(2,5-dimethoxyanilino)-2-phenyl-4-pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

- 25 -

4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-[3-(4-Bromo-1-methyl-1H-pyrazol-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(2-Methyl-pyrimidin-4-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Phenylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(3-Nitrophenyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(2,3,4,5,6-Pentafluorophenoxy)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(2-Ethyl-1-phenyl-pyrazolin-5-one-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(Phenylsulfone)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[3-(N-phenylamide)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 26 -

4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 27 -

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;

4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;

4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 28 -

4-(3-Methoxybenzylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Carboxyl-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-piperidino-anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 29 -

4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine;

4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Chloro-5-methoxyanilino)-2-morpholino-6-(methyl)pyrimidine;

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(2,5-Diethoxy-4-morpholinoanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

- 30 -

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide;

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-aminopyrimidine;

- 31 -

4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(N-Methyl-3-methoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methyl-4-nitroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(Indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Methoxy-2-methyl-4-nitroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3-Trifluoromethyl-1-pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(4,5-Dihydro-2-thiazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(1-Pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(4,5,6,7-Tetrahydro-indazol-1-yl)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(1H-3-Pyrazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-
pyrimidine;
6-Methyl-2-(2-pyridinyl)-4-(3-trifluoromethylbenzylamino)-pyrimidine;
6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine;
4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(4-Fluoro-3-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-
pyridinyl)pyrimidine;
4-(5-Chloro-2-methoxyanilino)-2-(3-pyridinyl)-6-
trifluoromethylpyrimidine;
4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(5-Hydroxy-2-nitroanilino)-2-(3-pyridinyl)-6-
trifluoromethylpyrimidine;
4-(2-Ethanesulfonyl-5-hydroxyanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-
trifluoromethylpyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(2-pyridinyl)-6-
trifluoromethylpyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-
trifluoromethylpyrimidine;

- 33 -

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Ethanesulfonyl-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethoxyphenylethylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3,4,5-Trichloroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3-Methoxy-dibenzofuran-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(1,5,6-Trimethyl-benzimidazol-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

- 34 -

4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(1,3,4-Triazol-1-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride;

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(*t*-butyl)pyrimidine;

4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine; and

4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine.

[0029] Useful alkyl groups include straight-chained and branched C₁₋₁₀ alkyl groups, more preferably C₁₋₆ alkyl groups. Typical C₁₋₁₀ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, 3-pentyl, hexyl and octyl groups, which may be optionally substituted.

- [0030] Useful alkoxy groups include oxygen substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted.
- [0031] Useful alkylthio groups include sulphur substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted. Also included are the sulfoxides and sulfones of such alkylthio groups.
- [0032] Useful amino groups include -NH₂, -NHR₁₅ and -NR₁₅R₁₆, wherein R₁₅ and R₁₆ are C₁₋₁₀ alkyl or cycloalkyl groups, or R₁₅ and R₁₆ are combined with the N to form a ring structure, such as a piperidine, or R₁₅ and R₁₆ are combined with the N and other group to form a ring, such as a piperazine. The alkyl group may be optionally substituted.
- [0033] Optional substituents on the alkyl and cycloalkyl groups include one or more halo, hydroxy, carboxyl, amino, nitro, cyano, C₁-C₆ acylamino, C₁-C₆ acyloxy, C₁-C₆ alkoxy, aryloxy, alkylthio, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, saturated and unsaturated heterocyclic or heteroaryl.
- [0034] Optional substituents on the aryl, arylalkyl and heteroaryl groups include one or more halo, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₁-C₆)alkyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, C₁-C₆ hydroxyalkyl, nitro, amino, ureido, cyano, C₁-C₆ acylamino, hydroxy, thiol, C₁-C₆ acyloxy, azido, C₁-C₆ alkoxy or carboxy.
- [0035] The term "aryl" as employed herein by itself or as part of another group refers to monocyclic, bicyclic or tricyclic aromatic groups containing from 6 to 14 carbons in the ring portion.
- [0036] Useful aryl groups include C₆₋₁₄ aryl, preferably C₆₋₁₀ aryl. Typical C₆₋₁₄ aryl groups include phenyl, naphthyl, phenanthrenyl, anthracenyl, indenyl, azulenyl, biphenyl, biphenylenyl and fluorenyl groups.
- [0037] Useful cycloalkyl groups are C₃₋₈ cycloalkyl. Typical cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

- 36 -

- [0038] Useful saturated or partially saturated carbocyclic groups are cycloalkyl groups as described above, as well as cycloalkenyl groups, such as cyclopentenyl, cycloheptenyl and cyclooctenyl.
- [0039] Useful halo or halogen groups include fluorine, chlorine, bromine and iodine.
- [0040] Useful arylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Preferably the arylalkyl group is benzyl, phenylethyl or naphthylmethyl.
- [0041] Useful haloalkyl groups include C₁₋₁₀ alkyl groups substituted by one or more fluorine, chlorine, bromine or iodine atoms, e.g., fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl, chloromethyl, chlorofluoromethyl and trichloromethyl groups.
- [0042] Useful acylamino (acylamido) groups are any C₁₋₆ acyl (alkanoyl) attached to an amino nitrogen, e.g., acetamido, chloroacetamido, propionamido, butanoylamido, pentanoylamido and hexanoylamido, as well as aryl-substituted C₁₋₆ acylamino groups, e.g., benzoylamido, and pentafluorobenzoylamido.
- [0043] Useful acyloxy groups are any C₁₋₆ acyl (alkanoyl) attached to an oxy (-O-) group, e.g., formyloxy, acetoxy, propionyloxy, butanoyloxy, pentanoyloxy and hexanoyloxy.
- [0044] The term heterocycle is used herein to mean a saturated or partially saturated 3-7 membered monocyclic, or 7-10 membered bicyclic ring system, which consists of carbon atoms and from one to four heteroatoms independently selected from the group consisting of O, N, and S, wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, the nitrogen can be optionally quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring, and wherein the heterocyclic ring can be substituted on carbon or on a nitrogen atom if the resulting compound is stable.
- [0045] Useful saturated or partially saturated heterocyclic groups include tetrahydrofuranyl, pyranal, piperidinyl, piperazinyl, pyrrolidinyl,

- 37 -

imidazolidinyl, imidazoliny, indoliny, isoindoliny, quinuclidiny, morpholiny, isochromanyl, chromanyl, pyrazolidiny, pyrazoliny, tetronoyl and tetramoyl groups.

[0046] The term "heteroaryl" as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 π electrons shared in a cyclic array; and containing carbon atoms and 1, 2 or 3 oxygen, nitrogen or sulfur heteroatoms.

[0047] Useful heteroaryl groups include thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, pyranal, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthinyl, 2*H*-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalziny, naphthyridiny, quinoxaliny, cinnoliny, pteridinyl, carbazolyl, β -carboliny, phenanthridiny, acridiny, perimidiny, phenanthroliny, phenaziny, isothiazolyl, phenothiaziny, isoxazolyl, furazanyl, phenoxaziny, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-a]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazolyl, 2-oxindolyl and 2-oxobenzimidazolyl. Where the heteroaryl group contains a nitrogen atom in a ring, such nitrogen atom may be in the form of an N-oxide, e.g., a pyridiny N-oxide, pyraziny N-oxide and pyrimidinyl N-oxide.


[0048] Some of the compounds of the present invention may exist as stereoisomers including optical isomers. The invention includes all stereoisomers and both the racemic mixtures of such stereoisomers as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art.

[0049] Examples of pharmaceutically acceptable addition salts include inorganic and organic acid addition salts such as hydrochloride, hydrobromide, phosphate, sulphate, citrate, lactate, tartrate, maleate, fumarate, mandelate and oxalate; and inorganic and organic base addition salts with bases such as sodium hydroxy, Tris(hydroxymethyl)aminomethane (TRIS, tromethane) and N-methyl-glucamine.

[0050] Examples of prodrugs of the compounds of the invention include the simple esters of carboxylic acid containing compounds (e.g., those obtained by condensation with a C₁₋₄ alcohol according to methods known in the art); esters of hydroxy containing compounds (e.g., those obtained by condensation with a C₁₋₄ carboxylic acid, C₃₋₆ dioic acid or anhydride thereof such as succinic and fumaric anhydrides according to methods known in the art); phosphate of hydroxy containing compounds (e.g. combretastatin A-1 phosphate prodrug, see Pettit G. R. and Lippert III, J. W., *Anticancer Drug Design* 15:203-216 (2000)); imines of amino containing compounds (e.g., those obtained by condensation with a C₁₋₄ aldehyde or ketone according to methods known in the art); carbamate of amino containing compounds such as those described by Leu, *et. al.*, (*J. Med. Chem.* 42:3623-3628 (1999)) and Greenwald, *et. al.*, (*J. Med. Chem.* 42:3657-3667 (1999)); and acetals and ketals of alcohol containing compounds (e.g., those obtained by condensation with chloromethyl methyl ether or chloromethyl ethyl ether according to methods known in the art).

[0051] The compounds of this invention may be prepared using methods known to those skilled in the art, or the novel methods of this invention. Specifically, the compounds of this invention with Formulae I-III may be prepared as illustrated by the exemplary reaction in Scheme 1. Reaction of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine with aniline gave the product 4-anilino-6-methyl-2-(2-pyridinyl)pyrimidine.

SCHEME 1




[0052] The 2-aryl-4-chloro-pyrimidine may be prepared as illustrated by the exemplary reaction in Scheme 2. Reaction of pyridine-2-carboxamide with

- 39 -

4,4,4-trifluoro-but-2-ynoic acid ethyl ester in ethanol in the presence of base such as KOH produced the substituted 4-hydroxy-pyrimidine. Treatment of the hydroxy-pyrimidine with POCl₃ gave the product 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine.


SCHEME 2



[0053] Alternatively, the 2-aryl-4-chloro-pyrimidine may be prepared as illustrated by the exemplary reaction in Scheme 3. Reaction of pyrimidine-2-carboxamide with ethyl 4,4,4-trifluoro-acetoacetate in ethanol in the presence of base such as EtONa produced the substituted 4-hydroxy-pyrimidine. Treatment of the hydroxy-pyrimidine with POCl₃ gave the product 4-chloro-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine.

- 40 -

SCHEME 3



[0054] An important aspect of the present invention is the discovery that compounds having Formula I-III are activators of caspases and inducers of apoptosis. Therefore, these compounds are useful in a variety of clinical conditions in which there is uncontrolled cell growth and spread of abnormal cells, such as in the case of cancer.

[0055] Another important aspect of the present invention is the discovery that compounds having Formula I-III are potent and highly efficacious activators of caspases and inducers of apoptosis in drug resistant cancer cells, such as breast cancer cells (Examples 11-14), which enables these compounds to kill these drug resistant cancer cells. In comparison, most standard anti-cancer drugs are not effective in killing drug resistant cancer cells under the same conditions. Therefore, compounds of this invention are useful for the treatment of drug resistant cancer such as breast cancer in animals.

[0056] The present invention includes a therapeutic method useful to modulate *in vivo* apoptosis or *in vivo* neoplastic disease, comprising administering to a subject in need of such treatment an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis.

[0057] The present invention also includes a therapeutic method comprising administering to an animal an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, wherein said therapeutic method is useful to treat cancer, which is a group of diseases characterized by the uncontrolled growth and spread of abnormal cells. Such diseases include, but are not limited to, Hodgkin's disease, non-Hodgkin's lymphomas, acute lymphatic leukemia, chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinomas, ovarian carcinomas, lung carcinomas, Wilms' tumor, cervical carcinomas, testicular carcinomas, soft-tissue sarcomas, primary macroglobulinemia, bladder carcinomas, chronic granulocytic leukemia, primary brain carcinomas, malignant melanoma, small-cell lung carcinomas, stomach carcinomas, colon carcinomas, malignant pancreatic insulinoma, malignant carcinoid carcinomas, malignant melanomas, choriocarcinomas, mycosis fungoides, head or neck carcinomas, osteogenic sarcoma, pancreatic carcinomas, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinomas, thyroid carcinomas, esophageal carcinomas, malignant hypercalcemia, cervical hyperplasia, renal cell carcinomas, endometrial carcinomas, polycythemia vera, essential thrombocytosis, adrenal cortex carcinomas, skin cancer, and prostatic carcinomas.

[0058] In practicing the therapeutic methods, effective amounts of compositions containing therapeutically effective concentrations of the compounds formulated for oral, intravenous, local and topical application, for the treatment of neoplastic diseases and other diseases in which caspase cascade mediated physiological responses are implicated, are administered to an individual exhibiting the symptoms of one or more of these disorders. The amounts are effective to ameliorate or eliminate one or more symptoms of the disorders. An effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce, the symptoms associated with the disease. Such amount may be administered as a

- 42 -

single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Typically, repeated administration is required to achieve the desired amelioration of symptoms

[0059] In another embodiment, a pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis in combination with a pharmaceutically acceptable vehicle is provided.

[0060] Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in combination with at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent. Examples of known cancer chemotherapeutic agents which may be used for combination therapy include, but not are limited to alkylating agents such as busulfan, cis-platin, mitomycin C, and carboplatin; antimitotic agents such as colchicine, vinblastine, paclitaxel, and docetaxel; topo I inhibitors such as camptothecin and topotecan; topo II inhibitors such as doxorubicin and etoposide; RNA/DNA antimetabolites such as 5-azacytidine, 5-fluorouracil and methotrexate; DNA antimetabolites such as 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea and thioguanine; antibodies such as Herceptin® and Rituxan®. Other known cancer chemotherapeutic agents which may be used for combination therapy include melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen and alanosine.

[0061] In practicing the methods of the present invention, the compound of the invention may be administered together with at least one known chemotherapeutic agent as part of a unitary pharmaceutical composition. Alternatively, the compound of the invention may be administered apart from

at least one known cancer chemotherapeutic agent. In one embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered substantially simultaneously, i.e. the compounds are administered at the same time or one after the other, so long as the compounds reach therapeutic levels *in vivo* at the same time. On another embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered according to their individual dose schedule, so long as the compounds reach therapeutic levels *in vivo*.

[0062] Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a bioconjugates of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in bioconjugation with at least one known therapeutically useful antibody, such as Herceptin® or Rituxan®, growth factors such as EGF, NGF, cytokines such as IL-2, IL-4, or any molecule that binds to the cell surface. The antibodies and other molecules will deliver the compound of Formulae I-III to its targets and make it an effective anticancer agent. The bioconjugates could also enhance the anticancer effect of therapeutically useful antibodies, such as Herceptin® or Rituxan®.

[0063] Similarly, another embodiment of the present invention is directed to a composition effective in inhibiting neoplasia comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in combination with radiation therapy. In this embodiment, the compound of the invention may be administered at the same time as the radiation therapy is administered or at a different time.

[0064] Yet another embodiment of the present invention is directed to a composition effective for post-surgical treatment of cancer, comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis. The invention also relates to a method of treating cancer by

- 44 -

surgically removing the cancer and then treating the animal with one of the pharmaceutical compositions described herein.

[0065] A wide range of immune mechanisms operate rapidly following exposure to an infectious agent. Depending on the type of infection, rapid clonal expansion of the T and B lymphocytes occurs to combat the infection. The elimination of the effector cells following an infection is one of the major mechanisms for maintaining immune homeostasis. The elimination of the effector cells has been shown to be regulated by apoptosis. Autoimmune diseases have lately been determined to occur as a consequence of deregulated cell death. In certain autoimmune diseases, the immune system directs its powerful cytotoxic effector mechanisms against specialized cells such as oligodendrocytes in multiple sclerosis, the beta cells of the pancreas in diabetes mellitus, and thyrocytes in Hashimoto's thyroiditis (Ohsako, S. & Elkon, K.B., *Cell Death Differ.* 6:13-21 (1999)). Mutations of the gene encoding the lymphocyte apoptosis receptor Fas/APO-1/CD95 are reported to be associated with defective lymphocyte apoptosis and autoimmune lymphoproliferative syndrome (ALPS), which is characterized by chronic, histologically benign splenomegaly, generalized lymphadenopathy, hypergammaglobulinemia, and autoantibody formation. (Infante, A.J., *et al.*, *J. Pediatr.* 133:629-633 (1998) and Vaishnav, A.K., *et al.*, *J. Clin. Invest.* 103:355-363 (1999)). It was reported that overexpression of Bcl-2, which is a member of the bcl-2 gene family of programmed cell death regulators with anti-apoptotic activity, in developing B cells of transgenic mice, in the presence of T cell dependent costimulatory signals, results in the generation of a modified B cell repertoire and in the production of pathogenic autoantibodies (Lopez-Hoyos, M., *et al.*, *Int. J. Mol. Med.* 1:475-483 (1998)). It is therefore evident that many types of autoimmune disease are caused by defects of the apoptotic process. One treatment strategy for such diseases is to turn on apoptosis in the lymphocytes that are causing the autoimmune disease (O'Reilly, L.A. & Strasser, A., *Inflamm. Res.* 48:5-21 (1999)).

- [0066] Fas-Fas ligand (FasL) interaction is known to be required for the maintenance of immune homeostasis. Experimental autoimmune thyroiditis (EAT), characterized by autoreactive T and B cell responses and a marked lymphocytic infiltration of the thyroid, is a good model to study the therapeutic effects of FasL. Batteux, F., *et al.*, (*J. Immunol.* 162:603-608 (1999)) reported that by direct injection of DNA expression vectors encoding FasL into the inflamed thyroid, the development of lymphocytic infiltration of the thyroid was inhibited and induction of infiltrating T cells death was observed. These results show that FasL expression on thyrocytes may have a curative effect on ongoing EAT by inducing death of pathogenic autoreactive infiltrating T lymphocytes.
- [0067] Bisindolylmaleimide VIII is known to potentiate Fas-mediated apoptosis in human astrocytoma 1321N1 cells and in Molt-4T cells; both of which were resistant to apoptosis induced by anti-Fas antibody in the absence of bisindolylmaleimide VIII. Potentiation of Fas-mediated apoptosis by bisindolylmaleimide VIII was reported to be selective for activated, rather than non-activated, T cells, and was Fas-dependent. Zhou T., *et al.*, (*Nat. Med.* 5:42-48 (1999)) reported that administration of bisindolylmaleimide VIII to rats during autoantigen stimulation prevented the development of symptoms of T cell-mediated autoimmune diseases in two models, the Lewis rat model of experimental allergic encephalitis and the Lewis adjuvant arthritis model. Therefore, the application of a Fas-dependent apoptosis enhancer such as bisindolylmaleimide VIII may be therapeutically useful for the more effective elimination of detrimental cells and inhibition of T cell-mediated autoimmune diseases. Therefore an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for autoimmune diseases.
- [0068] Psoriasis is a chronic skin disease that is characterized by scaly red patches. Psoralen plus ultraviolet A (PUVA) is a widely used and effective treatment for psoriasis vulgaris and Coven, *et al.*, *Photodermatol.*

Photoimmunol. Photomed. 15:22-27 (1999), reported that lymphocytes treated with psoralen 8-MOP or TMP and UVA, displayed DNA degradation patterns typical of apoptotic cell death. Ozawa, *et al.*, *J. Exp. Med.* 189:711-718 (1999) reported that induction of T cell apoptosis could be the main mechanism by which 312-nm UVB resolves psoriasis skin lesions. Low doses of methotrexate may be used to treat psoriasis to restore a clinically normal skin. Heenen, *et al.*, *Arch. Dermatol. Res.* 290:240-245 (1998), reported that low doses of methotrexate may induce apoptosis and that this mode of action could explain the reduction in epidermal hyperplasia during treatment of psoriasis with methotrexate. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for psoriasis.

[0069] Synovial cell hyperplasia is a characteristic of patients with rheumatoid arthritis (RA). It is believed that excessive proliferation of RA synovial cells, as well as defects in synovial cell death, may be responsible for synovial cell hyperplasia. Wakisaka, *et al.*, *Clin. Exp. Immunol.* 114:119-128 (1998), found that although RA synovial cells could die via apoptosis through a Fas/FasL pathway, apoptosis of synovial cells was inhibited by proinflammatory cytokines present within the synovium. Wakisaka, *et al.*, also suggested that inhibition of apoptosis by the proinflammatory cytokines may contribute to the outgrowth of synovial cells, and lead to pannus formation and the destruction of joints in patients with RA. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for rheumatoid arthritis.

[0070] There has been an accumulation of convincing evidence that apoptosis plays a major role in promoting resolution of the acute inflammatory response. Neutrophils are constitutively programmed to undergo apoptosis, thus limiting their pro-inflammatory potential and leading to rapid, specific, and non-phlogistic recognition by macrophages and semi-professional phagocytes

- 47 -

(Savill, J., *J. Leukoc. Biol.* 61:375-380 (1997)). Boirivant, *et al.*, *Gastroenterology* 116:557-565 (1999), reported that lamina propria T cells, isolated from areas of inflammation in Crohn's disease, ulcerative colitis, and other inflammatory states, manifest decreased CD2 pathway-induced apoptosis. In addition, studies of cells from inflamed Crohn's disease tissue indicate that this defect is accompanied by elevated Bcl-2 levels. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for inflammation.

[0071] Pharmaceutical compositions within the scope of this invention include all compositions wherein the compounds of the present invention are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the compounds may be administered to animals, e.g., mammals, orally at a dose of 0.0025 to 50 mg/kg of body weight, per day, or an equivalent amount of the pharmaceutically acceptable salt thereof, to a mammal being treated for apoptosis-mediated disorders. Preferably, about 0.01 to about 10 mg/kg of body weight is orally administered to treat or prevent such disorders. For intramuscular injection, the dose is generally about one-half of the oral dose. For example, a suitable intramuscular dose would be about 0.0025 to about 25 mg/kg of body weight, and most preferably, from about 0.01 to about 5 mg/kg of body weight. If a known cancer chemotherapeutic agent is also administered, it is administered in an amount that is effective to achieve its intended purpose. The amounts of such known cancer chemotherapeutic agents effective for cancer are well known to those of skill in the art.

[0072] The unit oral dose may comprise from about 0.01 to about 50 mg, preferably about 0.1 to about 10 mg of the compound of the invention. The unit dose may be administered one or more times daily as one or more tablets

- 48 -

each containing from about 0.1 to about 10, conveniently about 0.25 to 50 mg of the compound or its solvates.

[0073] In a topical formulation, the compound may be present at a concentration of about 0.01 to 100 mg per gram of carrier.

[0074] In addition to administering the compound as a raw chemical, the compounds of the invention may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the compounds into preparations which may be used pharmaceutically. Preferably, the preparations, particularly those preparations which may be administered orally and which may be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which may be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or orally, contain from about 0.01 to 99 percent, preferably from about 0.25 to 75 percent of active compound(s), together with the excipient.

[0075] Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like. Basic salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic base such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, Tris, N-methyl-glucamine and the like.

[0076] The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial effects of the compounds of the invention. Foremost among such animals are mammals,

- 49 -

e.g., humans and veterinary animals, although the invention is not intended to be so limited.

[0077] The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

[0078] The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use may be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

[0079] Suitable excipients are, in particular, fillers such as saccharides, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be

- 50 -

used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropymethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

[0080] Other pharmaceutical preparations which may be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules may contain the active compounds in the form of granules which may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

[0081] Possible pharmaceutical preparations which may be used rectally include, for example, suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example, liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

[0082] Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or

- 51 -

polyethylene glycol-400 (the compounds are soluble in PEG-400) or cremophor, or cyclodextrins. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

[0083] In accordance with one aspect of the present invention, compounds of the invention are employed in topical and parenteral formulations and are used for the treatment of skin cancer.

[0084] The topical compositions of this invention are formulated preferably as oils, creams, lotions, ointments and the like by choice of appropriate carriers. Suitable carriers include vegetable or mineral oils, white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high molecular weight alcohol (greater than C₁₂). The preferred carriers are those in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers may be employed in these topical formulations. Examples of such enhancers are found in U.S. Patent Nos. 3,989,816 and 4,444,762.

[0085] Creams are preferably formulated from a mixture of mineral oil, self-emulsifying beeswax and water in which mixture the active ingredient, dissolved in a small amount of an oil such as almond oil, is admixed. A typical example of such a cream is one which includes about 40 parts water, about 20 parts beeswax, about 40 parts mineral oil and about 1 part almond oil.

[0086] Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil such as almond oil with warm soft paraffin and allowing the mixture to cool. A typical example of such an ointment is one which includes about 30% almond oil and about 70% white soft paraffin by weight.

[0087] The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and

- 52 -

adaptations of the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

EXAMPLE 1

4-(4-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0088] To a mixture of 6-chloro-4-methyl-2-(2-pyridinyl)pyrimidine (115 mg, 0.56 mmol), *p*-anisidine (68 mg, 0.55 mmol) in water (10 mL) was added 2 N HCl (0.3 mL). The mixture was refluxed for 6 h, cooled to room temperature, and neutralized with 2N NaOH to pH 10. The resulting mixture was then extracted with 1:1 Hexane/EtOAc (50 mL). The organic phase was washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by chromatography (1:1 Hexane/EtOAc) to yield the product as a tan solid (72 mg, 45%). ¹H NMR (CDCl₃): 8.80 (d, *J* = 4.5 Hz, 1H), 8.45 (d, *J* = 8.1 Hz, 1H), 7.81 (m, 1H), 7.35 (m, 1H), 7.23 (d, *J* = 8.4 Hz, 2H), 7.12 (s, 1H), 6.93 (d, *J* = 8.4 Hz, 2H), 6.41 (s, 1H), 3.83 (s, 3H), 2.45 (s, 3H).

EXAMPLE 2

4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0089] The title compound was prepared using a similar method as described in Example 1. ¹H NMR (CDCl₃): 8.80 (d, *J* = 4.5 Hz, 1H), 8.46 (d, *J* = 7.5 Hz, 1H), 7.84 (m, 1H), 7.56 (s, 1H), 7.39-7.25 (m, 3H), 7.10 (d, *J* = 7.5 Hz, 1H), 6.83 (m, 1H), 6.64 (s, 1H), 2.45 (s, 3H).

- 53 -

EXAMPLE 3

4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

- [0090] The title compound was prepared using a similar method as described in Example 1. ^1H NMR (CDCl_3): 8.79 (d, $J = 4.5$ Hz, 1H), 8.44 (d, $J = 7.8$ Hz, 1H), 7.79 (m, 1H), 7.36-7.09 (m, 5H), 6.98 (d, $J = 7.5$ Hz, 1H), 6.59 (s, 1H), 2.46 (s, 3H), 2.35 (s, 3H).

EXAMPLE 4

4-Anilino-6-methyl-2-(2-pyridinyl)pyrimidine

- [0091] The title compound was prepared using a similar method as described in Example 1. ^1H NMR ($\text{DMSO}-d_6$): 9.64 (s, 1H), 8.70 (d, $J = 4.5$ Hz, 1H), 8.28 (d, $J = 7.5$ Hz, 1H), 7.94 (m, 1H), 7.78 (s, 1H), 7.75 (s, 1H), 7.47 (m, 1H), 7.31 (m, 1H), 7.02 (m, 1H), 6.66 (s, 1H), 2.38 (s, 3H).

EXAMPLE 5

4-(4-Methoxyanilino)-6-methyl-2-phenylpyrimidine

- [0092] The title compound was prepared using a similar method as described in Example 1. ^1H NMR (CDCl_3): 8.39-8.36 (m, 2H), 7.48-7.44 (m, 3H), 7.30-7.26 (m, 2H), 6.95-6.93 (m, 2H), 6.68 (s, 1H), 6.32 (s, 1H), 3.84 (s, 3H), 2.38 (s, 3H).

EXAMPLE 6

5-Chloro-4-(4-methoxyanilino)-6-methyl-2-[(5-trifluoromethyl)-2-pyridinyl]pyrimidine

- [0093] The title compound was prepared using a similar method as described in Example 1. ^1H NMR (CDCl_3): 9.06 (m, 1H), 8.43 (m, 1H), 8.04 (m, 1H), 7.59-7.56 (m, 2H), 7.23 (s, 1H), 7.00-6.97 (m, 2H), 3.87 (s, 3H), 2.71 (s, 3H).

EXAMPLE 7

6-(4-Chlorophenyl)-5-cyano-4-(4-methoxyanilino)-2-pyridinylpyrimidine

[0094] The title compound was prepared using a similar method as described in Example 1. ¹H NMR (DMSO-d₆): 8.58 (d, *J* = 4.5 Hz, 1H), 8.11 (d, *J* = 7.8 Hz, 1H), 7.88-7.77 (m, 3H), 7.54 (d, *J* = 8.7 Hz, 2H), 7.47-7.39 (m, 3H), 6.85 (d, *J* = 8.7 Hz, 2H), 3.64 (s, 3H).

EXAMPLE 8

4-(4-Methoxyanilino)-2-phenylquinazoline

[0095] To a solution of 4-chloro-2-phenylquinazoline (590 mg, 2.45 mmol) in THF (10 mL) was added *m*-anisidine (301 mg, 2.45 mmol) and NaH (60%, 120 mg, 3.0 mmol). The mixture was refluxed for 24 h. After cooled to room temperature, the mixture was diluted with 1:1 hexane/EtOAc (100 mL), washed with water and brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by chromatography (3:1 hexane/EtOAc) to yield the title compound as a white solid (173 mg, 0.53 mmol, 22%). ¹H NMR (CDCl₃): 8.54-8.51 (m, 2H), 8.00-7.74 (m, 5H), 7.52-7.47 (m, 4H), 7.38 (s, 1H), 7.02-7.00 (m, 2H), 3.88 (s, 3H).

EXAMPLE 9

6-Chloro-4-(4-methoxyanilino)-2-phenylpyrimidine

[0096] The title compound was prepared using a similar method as described in Example 8. ¹H NMR (CDCl₃): 8.39-8.36 (m, 1H), 8.48-7.46 (m, 3H), 7.28-7.25 (m, 3H), 6.98-6.95 (m, 2H), 6.85 (s, 1H), 6.42 (s, 1H), 3.85 (s, 3H).

- 55 -

EXAMPLE 10

4-(4-Chlorophenyl)-2-(4-methoxyanilino)-6-methylthio-1,3,5-triazine

[0097] The title compound was prepared using a similar method as described in Example 1. ^1H NMR (CDCl_3): 8.35 (d, $J = 8.4$ Hz, 1H), 7.52 (d, $J = 8.7$ Hz, 2H), 7.42 (d, $J = 8.7$ Hz, 2H), 7.16 (s, 1H), 6.92 (d, $J = 8.4$ Hz, 2H).

EXAMPLE 11

4-(4-Methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

[0098] A mixture of 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine (100 mg, 0.402 mmol), *p*-anisidine (50 mg, 0.402 mmol), water (5 ml) and 2N HCl (300 μl) was refluxed overnight. The resulting yellow solid was collected by filtration, washed with water and then dried under vacuum to give the title compound (55 mg, 41%). ^1H NMR (CDCl_3): 8.28 (d, $J = 8.7$ Hz, 2H), 7.43-7.35 (m, 4H), 6.96 (d, $J = 9.0$ Hz, 2H), 6.83 (s, 1H), 4.64 (s, 2H), 3.85 (s, 3H), 3.45 (s, 3H), 2.44 (s, 3H).

EXAMPLE 12

4-(4-Methoxyanilino)-6-methyl-2-(3-methylphenyl)pyrimidine

[0099] The title compound was prepared from 4-chloro-6-methyl-2-(3-methylphenyl)pyrimidine (100 mg, 0.457 mmol) and *p*-anisidine (56 mg, 0.457 mmol), similar to Example 11 as a pale green solid (39 mg, 28%). ^1H NMR ($\text{DMSO}-d_6$): 8.04 (s, 1H), 8.01-7.99 (m, 1H), 7.56-7.51 (m, 4H), 7.04 (d, $J = 9.0$ Hz, 2H), 6.73 (s, 1H), 3.79 (s, 3H), 2.44 (s, 6H).

EXAMPLE 13

4-(4-Dimethylaminoanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

[00100] A mixture of 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine (100 mg, 0.402 mmol), N,N-dimethyl-1,3-phenylene-diamine dihydrochloride (84 mg, 0.402 mmol), water (5 ml) and 2N HCl (300 μ l) was refluxed for 6 h. The mixture was diluted with ethyl acetate (50 ml), washed with water and saturated NaCl. The ethyl acetate solution was dried over anhydrous sodium sulfate, concentrated in vacuo and purified by column chromatography (hexane:ethyl acetate, 3:1) to yield a tan oily product (32 mg, 23%). ^1H NMR (CDCl_3): 8.22-8.20 (m, 1H), 8.18 (s, 1H), 7.34 (t, $J = 7.5$ Hz, 1H), 7.25- 7.20 (m, 2H), 6.96 (bs, 1H), 6.89 (s, 1H), 6.82 (s, 1H), 6.67 (dd, $J = 2.0, 8.0$ Hz, 1H), 6.53 (dd, $J = 2.6, 8.6$ Hz, 1H), 4.50 (m, 2H), 3.47 (m, 3H), 2.97 (s, 6H), 2.42 (s, 3H).

EXAMPLE 14

4-(4-Dimethylaminoanilino)-6-methyl-2-(3-methylphenyl)pyrimidine

[00101] The title compound was prepared from 4-chloro-6-methyl-2-(3-methylphenyl)pyrimidine (100 mg, 0.457 mmol) and N,N-dimethyl-1,3-phenylene-diamine dihydrochloride (96 mg, 0.457 mmol) similar to Example 13 as a yellow oil (9 mg, 6%). ^1H NMR (CDCl_3): 8.20 (s, 1H), 8.17 (s, 1H), 7.34 (t, $J = 7.5$ Hz, 1H), 7.25-7.21 (m, 2H), 6.82 (t, $J = 2.1$ Hz, 1H), 6.79 (bs, 1H), 6.66 (dd, $J = 2.0, 7.7$ Hz, 1H), 6.54 (dd, $J = 2.9, 8.9$ Hz, 2H), 2.97 (s, 6H), 2.42 (s, 6H).

- 57 -

EXAMPLE 15

4-(3-Methoxyanilino)-5-chloro-6-methyl-2-[5-(trifluoromethyl)-2-pyridinyl]pyrimidine

[0100] The title compound was prepared from 4,5-dichloro-6-methyl-2-[5-(trifluoromethyl)-2-pyridinyl]pyrimidine (100 mg, 0.325 mmol) and *m*-anisidine (37 μ l, 0.325 mmol) similar to Example 11 as a pale white solid (26 mg, 20%). ^1H NMR (CDCl_3): 9.05 (dd, $J = 0.8, 1.4$ Hz, 1H), 8.53 (d, $J = 8.4$ Hz, 1H), 8.06 (dd, $J = 2.3, 8.3$ Hz, 1H), 7.57 (t, $J = 2.3$ Hz, 1H), 7.36 (s, 1H), 7.31 (t, $J = 8.1$ Hz, 1H), 7.18-7.14 (m, 1H), 6.75-6.72 (m, 1H), 3.88 (s, 3H), 2.71 (s, 3H).

EXAMPLE 16

4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0101] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and *m*-anisidine (55 μ l, 0.486 mmol) similar to Example 13 as an tan oil (139 mg, 98%). ^1H NMR (CDCl_3): 8.78- 8.75 (m, 1H), 8.45 (dt, $J = 1.2, 8.1$ Hz, 1H), 7.78 (td, $J = 1.9, 7.8$ Hz, 1H), 7.73 (s, 1H), 7.34-7.30 (m, 1H), 7.23 (t, $J = 8.1$ Hz, 1H), 7.03 (t, $J = 2.1$ Hz, 1H), 6.92 (dd, $J = 2.0, 7.7$ Hz, 1H), 6.67 (dd, $J = 2.4, 8.1$ Hz, 1H), 6.61 (s, 1H), 3.77 (s, 3H), 2.44 (s, 3H).

EXAMPLE 17

4-(3-Methoxyanilino)-6-methyl-2-(3-methylphenyl)pyrimidine

[0102] The title compound was prepared from 4-chloro-6-methyl-2-(3-methylphenyl)pyrimidine (100 mg, 0.457 mmol) and *m*-anisidine (51 μ l, 0.457 mmol) similar to Example 11 as a pale yellow solid (11 mg, 8%). ^1H NMR ($\text{DMSO}-d_6$): 8.09 (s, 1H), 8.05 (d, $J = 7.5$ Hz, 1H), 7.54-7.49 (m, 4H), 7.35 (t, $J = 8.1$ Hz, 1H), 7.21 (s, 1H), 6.78 (s, 2H), 3.81 (s, 3H), 2.43 (s, 6H).

- 58 -

EXAMPLE 18

4-(3-Methoxyanilino)-5-methoxy-2-(2-pyridinyl)pyrimidine

- [0103] The title compound was prepared from 4-chloro-5-methoxy-2-(2-pyridinyl)pyrimidine (100 mg, 0.451 mmol) and *m*-anisidine (51 μ l, 0.451 mmol) similar to Example 13 and isolated as a tan oily liquid (63 mg, 45%). ^1H NMR (CDCl_3): 8.76-8.74 (m, 1H), 8.37 (dd, $J = 1.1, 8.0$ Hz, 1H), 8.05 (s, 1H), 7.79-7.73 (m, 2H), 6.65-6.62 (m, 1H), 3.92 (s, 3H), 3.84 (s, 3H).

EXAMPLE 19

4-(3-Methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0104] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (100 mg, 0.385 mmol) and *m*-anisidine (43 μ l, 0.385 mmol) similar to Example 13 and isolated as a pale yellow solid (38 mg, 29%). ^1H NMR (Acetone-d_6): 9.61 (d, $J = 2.4$ Hz, 1H), 9.42 (s, 1H), 8.76-8.70 (m, 2H), 7.57-7.53 (m, 2H), 7.39 (s, 1H), 7.36 (s, 1H), 7.15 (s, 1H), 6.79-6.75 (m, 1H), 3.87 (s, 3H).

EXAMPLE 20

4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

- [0105] The title compound was prepared from 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine (100 mg, 0.402 mmol) and *m*-anisidine (45 μ l, 0.402 mmol) similar to Example 13 and isolated as a tan solid (154 mg). ^1H NMR (CDCl_3): 8.23 (d, $J = 9.0$ Hz, 2H), 7.34 (s, 1H), 7.25-7.16 (m, 4H), 6.92-6.89 (m, 1H), 6.75 (s, 1H), 6.69-6.65 (m, 1H), 4.51 (s, 2H), 3.78 (s, 3H), 3.47 (s, 3H), 2.40 (s, 3H).

- 59 -

EXAMPLE 21

4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0106] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 2,5-dimethoxyaniline similar to Example 13 and isolated as a tan oil (116 mg, 74 %). ¹H NMR (CDCl₃): 8.78-8.76 (m, 1H), 8.44 (d, *J* = 7.8 Hz, 1H), 7.81 (d, *J* = 2.7 Hz, 1H), 7.79-7.74 (m, 1H), 7.32-7.28 (m, 2H), 6.78 (d, *J* = 9.3 Hz, 1H), 6.60 (s, 1H), 6.53 (dd, *J* = 3.0, 8.7 Hz, 1H), 3.77 (d, *J* = 3.3 Hz, 6H), 2.48 (s, 3H).

EXAMPLE 22

4-(2,3-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0107] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 2,3-dimethoxyaniline (69 μl, 0.486 mmol) similar to Example 11 and isolated as a white solid (169 mg). ¹H NMR (CDCl₃): 8.85-8.82 (m, 1H), 8.48 (d, *J* = 7.8 Hz, 1H), 7.87-7.82 (m, 1H), 7.54 (d, *J* = 7.5 Hz, 1H), 7.40-7.35 (m, 2H), 7.10 (t, *J* = 8.4 Hz, 1H), 6.73 (d, *J* = 1.5 Hz, 1H), 7.00 (s, 1H), 3.90 (s, 3H), 3.86 (s, 3H), 2.53 (s, 3H).

EXAMPLE 23

4-(3-Benzyloxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0108] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 3-benzyloxyaniline (97 mg, 0.486 mmol) similar to Example 13 and isolated as a yellow oil (161 mg, 90%). ¹H NMR (CDCl₃): 8.78-8.76 (m, 1H), 8.48-8.45 (m, 1H), 7.79-7.74 (m, 1H), 7.68 (s, 1H), 7.42-7.29 (m, 6H), 7.24 (t, *J* = 8.1 Hz, 1H), 7.10 (s, 1H), 6.92 (dd, *J* = 0.8, 8.0 Hz, 1H), 6.77-6.74 (m, 1H), 6.58 (s, 1H), 5.04 (s, 2H), 2.42 (s, 3H).

- 60 -

EXAMPLE 24

4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0109] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and *m*-toluidine (52 μ l, 0.486 mmol) similar to Example 13 and isolated as a pale yellow oil (132 mg, 99%). ^1H NMR (CDCl_3): 8.76-8.74 (m, 1H), 8.43-8.40 (m, 1H), 7.78-7.72 (m, 1H), 7.55 (s, 1H), 7.31-7.08 (m, 4H), 6.92 (d, $J = 7.2$ Hz, 1H), 6.55 (s, 1H), 2.41 (s, 3H), 2.30 (s, 3H).

EXAMPLE 25

4-(3-Ethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0110] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and *m*-phenetidine (65 μ l, 0.486 mmol) similar to Example 11 and isolated as a light tan solid (116 mg, 78%). ^1H NMR (CDCl_3): 8.86-8.83 (m, 1H), 8.50 (d, $J = 8.1$ Hz, 1H), 7.89-7.83 (m, 1H), 7.42-7.38 (m, 1H), 7.35-7.30 (m, 1H), 7.09 (s, 1H), 6.96-6.84 (m, 2H), 6.76 (d, $J = 3.3$ Hz, 1H), 6.71 (s, 1H), 4.09 (q, $J = 7.2$ Hz, 2H), 2.53 (s, 3H), 1.47 (t, $J = 7.1$ Hz, 3H).

EXAMPLE 26

4-(3-Cyanoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0111] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 3-aminobenzonitrile (57 mg, 0.486 mmol) similar to Example 11 and isolated as a pale white solid (107 mg, 76%). ^1H NMR ($\text{DMSO}-d_6$): 10.14 (s, 1H), 8.78-8.75 (m, 1H), 8.56 (s, 1H), 8.32 (d, $J = 7.8$ Hz, 1H), 8.04-7.96 (m, 2H), 7.60-7.47 (m, 3H), 6.75 (s, 1H), 2.47 (s, 3H).

- 61 -

EXAMPLE 27

4-(3-Acetophenoneanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

- [0112] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 3-aminobenzophenone (96 mg, 0.486 mmol) similar to Example 11 and isolated as a light yellow solid (137 mg, 77%). ¹H NMR (DMSO-d₆): 10.76 (s, 1H), 8.78 (d, *J* = 3.9 Hz, 1H), 8.31 (s, 1H), 8.22 (d, *J* = 8.7 Hz, 1H), 8.08 (d, *J* = 9.3 Hz, 1H), 8.02-7.97 (m, 1H), 7.81 (d, *J* = 7.5 Hz, 2H), 7.68-7.51 (m, 6H), 6.85 (s, 1H), 2.5 (s, 3H).

EXAMPLE 28

4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

- [0113] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (100 mg, 0.486 mmol) and 3-fluoroaniline (47 μl, 0.486 mmol) similar to Example 11 and isolated as a light brown solid (74 mg, 54%). ¹H NMR (CDCl₃): 8.83 (d, *J* = 4.8 Hz, 1H), 8.47 (d, *J* = 9 Hz, 1H), 7.88 (m, 1H), 7.44-7.30 (m, 4H), 7.18-7.14 (m, 1H), 6.92-6.86 (m, 1H), 6.80 (s, 1H), 2.55 (s, 3H).

EXAMPLE 29

4-(2-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

- [0114] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and *o*-anisidine (27 μl, 0.243 mmol) similar to Example 13 and isolated as a pale white solid (40 mg, 56 %). ¹H NMR (CDCl₃): 8.82-8.80 (m, 1H), 8.46 (d, *J* = 8.1 Hz, 1H), 7.90 (dd, *J* = 1.4, 7.7 Hz, 1H), 7.84-7.78 (m, 1H), 7.36-7.32 (m, 1H), 7.28 (s, 1H), 7.18-6.98 (m, 2H), 6.92 (dd, *J* = 1.8, 8.0 Hz, 1H), 6.63 (s, 1H), 3.85 (s, 3H), 2.50 (s, 3H).

EXAMPLE 30

4-(3-Trifluoromethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0115] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-aminobenzotrifluoride (30 μ l, 0.243 mmol) similar to Example 11 and isolated as a pale white solid (74 mg, 93%). ^1H NMR (CDCl_3): 8.84-8.82 (m, 1H), 8.49-8.45 (m, 1H), 7.95 (s, 1H), 7.90-7.85 (m, 1H), 7.62 (d, $J = 7.8$ Hz, 1H), 7.52 (t, $J = 7.8$ Hz, 1H), 7.45-7.40 (m, 2H), 6.78 (s, 1H), 2.55 (s, 3H).

EXAMPLE 31

4-(3-Acetoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0116] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-aminoacetophenone (33 mg, 0.243 mmol) similar to Example 11 and isolated as a white solid (60 mg, 81%). ^1H NMR (CDCl_3): 8.83 (d, $J = 4.5$ Hz, 1H), 8.50 (d, $J = 8.1$ Hz, 1H), 8.13 (s, 1H), 7.89-7.83 (m, 1H), 7.75 (d, $J = 7.8$ Hz, 1H), 7.61-7.48 (m, 2H), 7.41-7.37 (m, 1H), 7.12 (s, 1H), 6.61 (s, 1H), 2.65 (s, 3H), 2.53 (s, 3H).

EXAMPLE 32

4-(3-Trifluoromethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0117] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-(trifluoromethoxy)aniline (65 μ l, 0.243 mmol) similar to Example 11 and isolated as a pale white solid (78 mg, 93%). ^1H NMR ($\text{DMSO}-d_6$): 9.95 (s, 1H), 8.74 (s, 1H), 8.31 (d, $J = 9.3$ Hz, 2H), 7.95 (t, $J = 7.7$ Hz, 1H), 7.65 (d, $J = 8.4$ Hz, 1H), 7.53-7.43 (m, 2H), 6.98 (d, $J = 8.4$ Hz, 1H), 6.71 (s, 1H), 2.44 (s, 3H).

EXAMPLE 33

4-(3-Ethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0118] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-ethylaniline (60 μ l, 0.243 mmol) similar to Example 13 and isolated as a tan oil (69 mg, 98%). ^1H NMR (CDCl_3): 8.81-8.79 (m, 1H), 8.48-8.45 (m, 1H), 7.84-7.78 (m, 1H), 7.42 (s, 1H), 7.37-7.29 (m, 2H), 7.18 (s, 1H), 7.16 (d, $J = 1.5$ Hz, 1H), 7.02 (d, $J = 7.8$ Hz, 1H), 6.61 (s, 1H), 2.66 (q, $J = 7.5$ Hz, 2H), 2.47 (s, 3H), 1.25 (t, $J = 7.5$ Hz, 3H).

EXAMPLE 34

4-(3-Nitroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0119] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-nitroaniline (34 mg, 0.243 mmol) similar to Example 11 as a yellow solid (65 mg, 87%). ^1H NMR (DMSO-d_6): 10.17 (s, 1H), 9.18 (s, 1H), 8.77 (d, $J = 4.8$ Hz, 1H), 8.39 (d, $J = 8.1$ Hz, 1H), 8.13 (d, $J = 7.2$ Hz, 1H), 7.98 (t, $J = 6.9$ Hz, 1H), 7.86 (d, $J = 8.4$ Hz, 1H), 7.63 (t, $J = 8.3$ Hz, 1H), 7.55-7.51 (m, 1H), 6.74 (s, 1H), 2.47 (s, 3H).

EXAMPLE 35

4-(3-Methylthioanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0120] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-(methylthio)aniline (30 μ l, 0.243 mmol) similar to Example 13 and isolated as a pink oil (67 mg, 89%). ^1H NMR (CDCl_3): 8.77-8.74 (m, 1H), 8.44-8.41 (m, 1H), 7.81-7.75 (m, 1H), 7.46 (s, 1H), 7.34-7.28 (m, 2H), 7.23 (t, $J = 8.1$ Hz, 1H), 7.09-7.06 (m, 1H), 7.00-6.97 (m, 1H), 6.55 (d, $J = 0.6$ Hz, 1H), 2.44 (s, 3H), 2.43 (d, $J = 0.6$, 3H).

- 64 -

EXAMPLE 36

4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0121] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and N,N-dimethyl-m-phenylenediamine (51 mg, 0.243 mmol) similar to Example 13 and isolated as a dark green oil (49 mg, 66%). ¹H NMR (CDCl₃): 8.82-8.79 (m, 1H), 8.47 (d, *J* = 7.8 Hz, 1H), 7.84-7.78 (m, 1H), 7.37-7.21 (m, 3H), 6.68-6.66 (m, 3H), 6.58-6.54 (m, 1H), 2.96 (s, 6H), 2.46 (s, 3H).

EXAMPLE 37

4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0122] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-isopropylaniline (33 μl, 0.243 mmol) similar to Example 13 and isolated as a gray oil (70 mg, 95%). ¹H NMR (CDCl₃): 8.79-8.77 (m, 1H), 8.47-8.43 (m, 1H), 7.81-7.75 (m, 1H), 7.35-7.25 (m, 3H), 7.17-7.12 (m, 2H), 7.02 (d, *J* = 7.5 Hz, 1H), 2.93-2.84 (m, 1H), 2.45 (s, 3H), 1.24 (d, *J* = 6.9 Hz, 6H).

EXAMPLE 38

4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0123] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (100 mg, 0.385 mmol) and 2,5-dimethoxyaniline (59 mg, 0.385 mmol) similar to Example 13 and isolated as a greenish-yellow solid (24 mg, 17%). ¹H NMR (CDCl₃): 9.67-9.66 (m, 1H), 8.74-8.70 (m, 2H), 8.02 (s, 1H), 7.44-7.39 (m, 1H), 6.91 (s, 1H), 7.67 (s, 1H), 6.87 (d, *J* = 8.7 Hz, 1H), 6.65 (dd, *J* = 3.0, 9.0 Hz, 1H), 3.86 (d, *J* = 7.2 Hz, 6H).

- 65 -

EXAMPLE 39

4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

- [0124] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl) pyrimidine (100 mg, 0.402 mmol) and 2,5-dimethoxyaniline (62 mg, 0.402 mmol) similar to Example 13 and isolated as a pale white solid (65 mg, 45%). ¹H NMR (CDCl₃): 8.30 (d, *J* = 3.0 Hz, 1H), 8.26-8.23 (m, 2H), 7.37-7.25 (m, 3H), 6.81 (d, *J* = 9.0 Hz, 1H), 6.73 (s, 1H), 6.55 (dd, *J* = 3.0, 8.7 Hz, 1H), 4.52 (d, *J* = 0.9 Hz, 2H), 3.87 (s, 3H), 3.83 (s, 3H), 3.51 (s, 3H), 2.42 (s, 3H).

EXAMPLE 40

4-(3-Methoxy-phenoxy)-6-methyl-2-(2-pyridinyl)pyrimidine

- [0125] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3-methoxyphenol (27 μ l, 0.243 mmol) similar to Example 13 and isolated as a white oil (24mg, 34%). ¹H NMR (CDCl₃): 8.79-8.76 (m, 1H), 8.26-8.23 (m, 1H), 7.75-7.69 (m, 1H), 7.33-7.28 (m, 2H), 6.82-6.72 (m, 3H), 6.58 (s, 1H), 3.77 (s, 3H), 2.58 (s, 3H).

EXAMPLE 41

4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0126] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and *m*-anisidine (22 μ l, 0.193 mmol) similar to Example 13 and isolated as a yellow solid (31mg, 46%). ¹H NMR (CDCl₃): 8.77-8.75 (m, 2H), 8.28-8.25 (m, 2H), 7.36 (t, *J* = 8.3 Hz, 1H), 7.13 (s, 1H), 7.03 (s, 1H), 6.98-6.96 (m, 2H), 6.83-6.81 (m, 1H), 3.84 (s, 3H).

- 66 -

EXAMPLE 42

4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0127] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2,5-dimethoxyaniline (32 mg, 0.193 mmol) similar to Example 13 and isolated as a light tan solid (15 mg, 21%). ¹H NMR (CDCl₃): 8.79 (dd, *J* = 1.8, 4.8 Hz, 2H), 8.31 (dd, *J* = 1.8, 4.8 Hz, 2H), 8.02 (bs, 1H), 7.59 (s, 1H), 6.98 (s, 1H), 6.90 (d, *J* = 9.0 Hz, 1H), 6.69 (dd, *J* = 2.9, 9.2 Hz, 1H), 3.90 (s, 3H), 3.87 (s, 3H).

EXAMPLE 43

4-(2,5-Dimethoxyanilino)-5-methoxy-2-(2-pyridinyl)pyrimidine

[0128] The title compound was prepared from a mixture of 4-chloro-5-methoxy-2-(2-pyridinyl)pyrimidine (50 mg, 0.226 mmol) and 2,5-dimethoxyaniline (35 mg, 0.226 mmol) similar to Example 11 and isolated as a gray solid (46 mg, 61%). ¹H NMR (CDCl₃): 8.77-8.74 (m, 1H), 8.72 (d, *J* = 3.0 Hz, 1H), 8.43 (d, *J* = 8.1 Hz, 1H), 8.10 (s, 1H), 8.03 (s, 1H), 7.83-7.77 (m, 1H), 7.33-7.29 (m, 1H), 6.84 (d, *J* = 8.7 Hz, 1H), 6.56 (dd, *J* = 3.0, 8.7 Hz, 1H), 4.04 (s, 3H), 3.90 (s, 3H), 3.89 (s, 3H).

EXAMPLE 44

4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1-3-thiazol-4-yl)pyrimidine

[0129] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(2-methyl-1-3-thiazol-4-yl)pyrimidine (50 mg, 0.196 mmol) and m-anisidine (22 μl, 0.196 mmol) similar to Example 11 and isolated as a white solid (29 mg, 43%). ¹H NMR (CDCl₃): 8.10 (s, 1H), 7.28 (t, *J* = 8.1 Hz, 1H), 6.97 (s, 2H), 6.90 (d, *J* = 7.5 Hz, 1H), 6.85 (s, 1H), 6.73-6.70 (m, 1H), 4.52 (s, 2H), 3.82 (s, 3H), 3.45 (s, 3H), 2.81 (s, 3H).

- 67 -

EXAMPLE 45

4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1-3-thiazol-4-yl)pyrimidine

[0130] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(2-methyl-1-3-thiazol-4-yl)pyrimidine (50 mg, 0.196 mmol) and 2,5-dimethoxyaniline (33 mg, 0.196 mmol) similar to Example 11 and isolated as a gray solid (33 mg, 45%). ¹H NMR (CDCl₃): 8.11 (s, 1H), 7.96 (s, 1H), 7.36 (s, 1H), 6.84-6.81 (m, 2H), 6.56 (dd, *J* = 3.0, 9.0 Hz, 1H), 4.56 (s, 2H), 3.83 (s, 3H), 3.82 (s, 3H), 3.48 (s, 3H), 2.80 (s, 3H).

EXAMPLE 46

6-Morpholino-4-(3-methoxyanilino)-2-phenyl-pyrimidine

[0131] A mixture of 4-(6-chloro-2-phenyl-4-pyrimidyl)morpholine (50 mg, 0.181 mmol), m-anisidine (20 μl, 0.181 mmol) and aqueous 2N HCl (150 μl) in water:ethanol (1:1, 10 ml) was refluxed for 5 days. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (50 ml), which was washed with water (2x25 ml), aqueous saturated NaCl, and dried over anhydrous sodium sulfate. The ethyl acetate solution was concentrated in vacuo and isolated as a yellow solid (6 mg, 9%). ¹H NMR (CDCl₃): 8.40-8.37 (m, 2H), 7.46-7.43 (m, 3H), 7.31-7.25 (m, 2H), 7.00 (t, *J* = 2.3 Hz, 1H), 6.91 (dd, *J* = 1.7, 8.0 Hz, 1H), 6.70-6.67 (m, 2H), 3.83 (s, 3H), 3.81-3.63 (m, 8H).

EXAMPLE 47

6-Morpholino-4-(2,5-dimethoxyanilino)-2-phenyl-4-pyrimidine

[0132] The title compound was prepared from a mixture of 4-(6-chloro-2-phenyl-4-pyrimidyl)morpholine (50 mg, 0.181 mmol) and 2,5-dimethoxyaniline (28 mg, 0.181 mmol) similar to Example 46 and isolated as a brown oil (6 mg, 8%). ¹H NMR (DMSO-d₆): 8.49 (bs, 1H), 8.33-8.30 (m,

- 68 -

2H), 7.93 (bs, 1H), 6.22 (s, 1H), 7.52-7.50 (m, 3H), 6.99 (dd, $J = 8.7$ Hz, 1H), 6.61 (dd, $J = 2.7, 8.7$ Hz, 1H), 3.81 (s, 3H), 3.76 (s, 3H), 3.74-3.70 (m, 8H).

EXAMPLE 48

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine

[0133] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine (50 mg, 0.196 mmol) and 2-chloro-5-methoxyaniline (38 mg, 0.196 mmol) similar to Example 13 and isolated as a white solid (33 mg, 45%). ^1H NMR (CDCl_3): 8.14 (s, 1H), 7.93 (d, $J = 3.0$ Hz, 1H), 7.32 (d, $J = 8.7$ Hz, 1H), 7.22 (s, 1H), 6.86 (s, 1H), 6.64 (dd, $J = 3.0, 9.3$ Hz, 1H), 4.60 (s, 2H), 3.88 (s, 3H), 3.52 (s, 3H), 2.84 (s, 3H).

EXAMPLE 49

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine

[0134] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine (50 mg, 0.196 mmol) and 5-methoxy-2-methylaniline (27 mg, 0.196 mmol) similar to Example 13 and isolated as a tan oil (42 mg, 60%). ^1H NMR (CDCl_3): 8.12 (s, 1H), 7.19 (d, $J = 8.4$ Hz, 1H), 6.99 (d, $J = 2.4$ Hz, 1H), 6.89 (s, 1H), 6.75 (dd, $J = 3.0, 8.7$ Hz, 1H), 6.56 (s, 1H), 4.50 (s, 2H), 3.80 (s, 3H), 3.44 (s, 3H), 2.83 (s, 3H), 2.21 (s, 3H).

EXAMPLE 50

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0135] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-chloro-5-

- 69 -

methoxyaniline (37 mg, 0.193 mmol) similar to Example 13 and isolated as a white solid (10 mg, 14%). ¹H NMR (CDCl₃): 9.70 (d, *J* = 1.8 Hz, 1H), 8.77-8.74 (m, 2H), 7.86 (s, 1H), 7.67 (s, 1H), 7.48-7.44 (m, 1H), 7.42 (s, 1H), 7.39 (s, 1H), 6.96 (s, 1H), 6.76 (dd, *J* = 3.0, 9.3 Hz, 1H), 3.90 (s, 3H).

EXAMPLE 51

4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0136] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-methoxy-2-methylaniline (26 mg, 0.193 mmol) similar to Example 13 and isolated as a tan solid (8 mg, 12%). ¹H NMR (CDCl₃): 9.72 (d, *J* = 0.6 Hz, 1H), 8.74-8.68 (m, 2H), 7.70 (s, 1H), 7.44-7.40 (m, 1H), 7.27 (s, 1H), 7.25 (s, 1H), 6.99 (s, 1H), 6.85 (dd, *J* = 2.7, 8.7 Hz, 1H), 6.64 (s, 1H), 3.83 (s, 3H), 2.24 (s, 3H).

EXAMPLE 52

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0137] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 2-chloro-5-methoxyaniline (47 mg, 0.243 mmol) similar to Example 13 and isolated as a tan oil (35 mg, 44%). ¹H NMR (CDCl₃): 8.82-8.80 (m, 1H), 8.47 (d, *J* = 7.8 Hz, 1H), 7.86-7.81 (m, 1H), 7.77 (d, *J* = 2.7 Hz, 1H), 7.41-7.36 (m, 1H), 7.36 (s, 1H), 7.30 (s, 1H), 6.68 (s, 1H), 6.65 (dd, *J* = 2.7, 8.7 Hz, 1H), 3.85 (s, 3H), 2.55 (s, 3H).

- 70 -

EXAMPLE 53

4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0138] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 5-methoxy-2-methylaniline (33 mg, 0.243 mmol) similar to Example 13 and isolated as a tan oil (45 mg, 60%). ¹H NMR (CDCl₃): 8.78-8.77 (m, 1H), 8.44 (d, *J* = 8.1 Hz, 1H), 7.81-7.75 (m, 1H), 7.34-7.30 (m, 1H), 7.16 (d, *J* = 8.4 Hz, 1H), 6.91 (d, *J* = 2.7 Hz, 2H), 6.72 (dd, *J* = 2.7, 8.7 Hz, 1H), 6.30 (s, 1H), 3.76 (s, 3H), 2.42 (s, 3H), 2.17 (s, 3H).

EXAMPLE 54

4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0139] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-chloro-5-methoxyaniline (38 mg, 0.193 mmol) similar to Example 13 and isolated as a pink solid (18 mg, 25%). ¹H NMR (CDCl₃): 8.82 (dd, *J* = 1.4, 4.7 Hz, 2H), 8.32 (dd, *J* = 1.7, 4.4 Hz, 2H), 7.84 (d, *J* = 2.1 Hz, 1H), 7.50 (s, 1H), 7.41 (d, *J* = 8.7 Hz, 1H), 7.02 (s, 1H), 6.77 (dd, *J* = 3.3, 9.3 Hz, 1H), 3.90 (s, 3H).

EXAMPLE 55

4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0140] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-methoxy-2-methylaniline (27 mg, 0.193 mmol) similar to Example 13 and isolated as a pink solid (23 mg, 33%). ¹H NMR (CDCl₃): 8.77 (d, *J* = 5.7 Hz, 2H), 8.27 (d, *J* = 5.7 Hz, 2H), 7.32 (s, 1H), 7.26 (d, *J* = 9.0 Hz, 1H), 6.99 (s, 1H), 6.70 (s, 1H), 6.85 (dd, *J* = 2.7, 8.4 Hz, 1H), 3.82 (s, 3H), 2.24 (s, 3H).

EXAMPLE 56

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

[0141] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine (50 mg, 0.201 mmol) and 2-chloro-5-methoxyaniline (39 mg, 0.201 mmol) similar to Example 13 and isolated as a white solid (18 mg, 24%). ¹H NMR (CDCl₃): 8.27-8.25 (m, 3H), 7.40 (t, *J* = 7.7 Hz, 1H), 7.34 (d, *J* = 8.7 Hz, 1H), 7.33 (s, 1H), 7.19 (s, 1H), 6.83 (s, 1H), 6.66 (dd, *J* = 3.0, 9.0 Hz, 1H), 4.59 (s, 2H), 3.93 (s, 3H), 3.58 (s, 3H), 2.47 (s, 3H).

EXAMPLE 57

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine

[0142] The title compound was prepared from a mixture of 4-chloro-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine (50 mg, 0.201 mmol) and 5-methoxy-2-methylaniline (28 mg, 0.201 mmol) similar to Example 13. ¹H NMR (CDCl₃): 8.20-8.16 (m, 2H), 7.35 (t, *J* = 7.5 Hz, 1H), 7.28 (s, 1H), 7.18 (s, 1H), 7.17 (d, *J* = 8.1 Hz, 1H), 6.73 (dd, *J* = 2.7, 8.4 Hz, 1H), 6.61 (s, 1H), 6.57 (s, 1H), 4.49 (s, 2H), 3.82 (s, 3H), 3.47 (s, 3H), 2.44 (s, 3H), 2.23 (s, 3H).

EXAMPLE 58

4-[3-(4-Bromo-1-methyl-1H-pyrazol-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0143] A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 3-(4-bromo-1-methyl-1H-pyrazol-3-yl)aniline (97 mg, 0.386 mmol) and aqueous 2N HCl (150 μl) in water:ethanol (1:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature, extracted with ethyl acetate (50 ml) and washed with water (1x25 ml). The ethyl acetate

- 72 -

solution was acidified with aqueous 2N HCl (1x25 ml). The resulting precipitate was collected by filtration, washed with water and isolated as a pale white solid (59 mg, 64%). ¹H NMR (DMSO-d₆): 10.54 (s, 1H), 9.44 (d, *J* = 2.1 Hz, 1H), 8.75 (dd, *J* = 1.8, 4.8 Hz, 1H), 8.62-8.58 (m, 1H), 7.94 (s, 1H), 7.92 (s, 1H), 7.71 (s, 1H), 7.65 (t, *J* = 8.3 Hz, 1H), 7.59-7.54 (m, 1H), 7.28 (d, *J* = 8.1 Hz, 1H), 7.19 (s, 1H), 3.81 (s, 3H).

EXAMPLE 59

4-[3-(2-Methyl-pyrimidin-4-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0144] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-(3-aminophenyl)-2-methylpyrimidine (71 mg, 0.386 mmol) similar to Example 58. The mixture was extracted with ethyl acetate (50 ml), washed with water (1x25 ml) and with aqueous 2N HCl (1x25 ml). The acidic aqueous solution was basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water and isolated as a light tan solid (18 mg, 23%). ¹H NMR (CDCl₃): 9.71 (d, *J* = 2.4 Hz, 1H), 8.77-8.72 (m, 3H), 8.38 (s, 1H), 7.89 (d, *J* = 7.2 Hz, 1H), 7.69-7.54 (m, 4H), 7.45-7.41 (m, 1H), 6.95 (s, 1H), 2.83 (s, 3H).

EXAMPLE 60

4-(3-Phenylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0145] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-aminobiphenyl (65 mg, 0.386 mmol) similar to Example 58. The mixture was basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was collected by filtration, washed with excess water:ethanol (1:1) and isolated as a light brown solid (64 mg, 85%). ¹H NMR (CDCl₃): 9.67 (s, 1H), 8.75 (d, *J* = 5.4 Hz, 2H), 7.68 (s, 1H), 7.63-7.37 (m, 10H), 6.98 (s, 1H).

- 73 -

EXAMPLE 61

4-[3-(3-Nitrophenyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0146] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-amino-3'-nitrobiphenyl (83 mg, 0.386 mmol) similar to Example 58 and isolated as a light tan solid (65 mg, 77%). ¹H NMR (DMSO-d₆): 13.00 (s, 1H), 11.94 (s, 1H), 11.23 (d, *J* = 4.5 Hz, 1H), 11.11 (d, *J* = 8.1 Hz, 1H), 10.97 (s, 1H), 10.78 (s, 1H), 10.75 (s, 1H), 10.69 (d, *J* = 8.1 Hz, 1H), 10.35-10.28 (m, 2H), 10.14-10.04 (m, 3H), 9.67 (s, 1H).

EXAMPLE 62

4-[3-(2,3,4,5,6-Pentafluorophenoxy)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0147] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-(2,3,4,5,6-pentafluorophenoxy)aniline (106 mg, 0.386 mmol) similar to Example 58 and isolated as a yellow solid (65 mg, 68%). ¹H NMR (DMSO-d₆): 10.46 (s, 1H), 9.24 (s, 1H), 8.75 (dd, *J* = 1.8, 4.8 Hz, 1H), 8.56-8.52 (m, 1H), 7.77 (s, 1H), 7.59-7.55 (m, 1H), 7.50-7.41 (m, 2H), 7.14 (s, 1H), 6.94-6.92 (m, 1H).

EXAMPLE 63

4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0148] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2,5-dimethoxyaniline (59 mg, 0.386 mmol) similar to Example 58 and isolated as a light brown solid (36 mg, 50%). ¹H NMR (DMSO-d₆): 9.65 (s, 1H), 8.76-8.73 (m, 1H), 8.32 (d, *J* = 7.8 Hz, 2H), 8.03-7.97 (m, 1H), 7.58-7.54 (m, 2H),

- 74 -

7.05 (d, $J = 9.0$ Hz, 1H), 6.70 (dd, $J = 3.0, 8.7$ Hz, 1H), 3.84 (s, 3H), 3.82 (s, 3H).

EXAMPLE 64

4-[3-(2-Ethyl-1-phenyl-pyrazolin-5-one-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0149] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-(3-aminophenyl)-2-ethyl-1-phenyl-3-pyrazolin-5-one (54 mg, 0.193 mmol) similar to Example 58 and isolated as a dark yellow solid (52 mg, 54%). ^1H NMR (DMSO- d_6): 10.61 (s, 1H), 9.52 (s, 1H), 8.83-8.81 (m, 1H), 8.65 (dd, $J = 1.2, 7.5$ Hz, 1H), 8.38 (s, 1H), 7.80 (d, $J = 7.8$ Hz, 1H), 7.68-7.48 (m, 7H), 7.37-7.33 (m, 1H), 7.22 (s, 1H), 5.97 (s, 1H), 3.63 (q, $J = 6.6$ Hz, 2H), 0.634 (t, $J = 6.6$ Hz, 3H).

EXAMPLE 65

4-[3-(Phenylsulfone)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0150] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-aminodiphenylsulfone (90 mg, 0.386 mmol) similar to Example 58 and isolated as a yellow solid (44 mg, 50%). ^1H NMR (CDCl_3): 9.68 (d, $J = 2.4$ Hz, 1H), 9.16 (s, 1H), 8.78-8.74 (m, 1H), 8.71-8.69 (m, 1H), 8.61 (s, 1H), 7.97 (d, $J = 7.2$ Hz, 2H), 7.85-7.82 (m, 1H), 7.65 (d, $J = 7.5$ Hz, 1H), 7.56-7.42 (m, 5H), 6.94 (s, 1H).

EXAMPLE 66

4-[3-(N-phenylamide)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0151] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-

- 75 -

aminobenzanilide (41 mg, 0.193 mmol) similar to Example 58 and isolated as a dark pink solid (41 mg, 49%). ¹H NMR (DMSO-d₆): 10.43 (s, 1H), 10.24 (s, 1H), 9.39 (d, *J* = 1.2 Hz, 1H), 8.62 (dd, *J* = 1.5, 4.8 Hz, 1H), 8.57-8.54 (m, 1H), 8.40 (s, 1H), 7.85 (d, *J* = 7.2 Hz, 1H), 7.69 (d, *J* = 8.4 Hz, 2H), 7.63 (d, *J* = 7.2 Hz, 1H), 7.50 (t, *J* = 8.0 Hz, 1H), 7.43-7.39 (m, 1H), 7.25 (t, *J* = 7.8 Hz, 2H), 7.06 (s, 1H), 7.02-6.97 (m, 1H).

EXAMPLE 67

4-(3-Phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0152] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-phenoxyaniline (71 mg, 0.386 mmol) similar to Example 58 and isolated as a pale white solid (33 mg, 42%). ¹H NMR (CDCl₃): 9.74 (d, *J* = 2.1 Hz, 1H), 8.72-8.66 (m, 2H), 8.52 (bs, 1H), 7.45-7.37 (m, 4H), 7.21-7.10 (m, 5H), 6.97 (s, 1H), 6.93 (dd, *J* = 2.1, 8.1 Hz, 1H).

EXAMPLE 68

4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0153] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2,5-dimethylaniline (48 μl, 0.386 mmol) similar to Example 58 and isolated as a light tan solid (14 mg, 21%). ¹H NMR (CDCl₃): 9.70 (dd, *J* = 0.8, 2.0 Hz, 1H), 8.74-8.70 (m, 1H), 8.67 (dd, *J* = 1.8, 4.8 Hz, 1H), 7.75 (s, 1H), 7.43-7.38 (m, 1H), 7.24 (d, *J* = 7.5 Hz, 1H), 7.19 (s, 1H), 7.10 (d, *J* = 7.8 Hz, 1H), 6.56 (s, 1H), 2.38 (s, 3H), 2.26 (s, 3H).

- 76 -

EXAMPLE 69

4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride

[0154] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (89 mg, 0.344 mmol) and 3-aminophenol (30 mg, 0.275 mmol) similar to Example 58 and isolated as a yellow solid (61 mg, 67%). ¹H NMR (DMSO-d₆): 10.40 (s, 1H), 8.86 (dd, *J* = 0.9, 5.1 Hz, 1H), 8.51 (d, *J* = 7.8 Hz, 1H), 8.32-8.26 (m, 1H), 7.82-7.78 (m, 1H), 7.29 (s, 1H), 7.26 (s, 1H), 7.23 (s, 1H), 7.19 (d, *J* = 7.5 Hz, 1H), 6.57 (dd, *J* = 1.8, 9.0 Hz, 1H).

EXAMPLE 70

4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0155] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,4-methylenedioxyaniline (53 mg, 0.386 mmol) similar to Example 58 and isolated as a purple solid (24 mg, 35%). ¹H NMR (CDCl₃): 8.85-8.83 (m, 1H), 8.57-8.54 (m, 1H), 7.90-7.84 (m, 1H), 7.45-7.41 (m, 1H), 7.39 (s, 1H), 6.87 (d, *J* = 8.1 Hz, 1H), 6.84 (s, 1H), 6.80 (s, 1H), 6.77 (dd, *J* = 2.1, 8.1 Hz, 1H), 6.05 (s, 2H).

EXAMPLE 71

4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0156] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,4-methylenedioxyaniline (53 mg, 0.386 mmol) similar to Example 58 and isolated as a dark brown solid (47 mg, 68%). ¹H NMR (CDCl₃): 8.77 (dd, *J* = 1.5, 4.5 Hz, 2H), 8.26 (dd, *J* = 1.5, 4.5 Hz, 2H), 7.06 (s, 1H), 6.93 (s, 1H),

- 77 -

6.89 (d, $J = 8.1$ Hz, 1H), 6.81 (dd, $J = 2.1, 8.1$ Hz, 1H), 6.81 (s, 1H), 6.06 (s, 2H).

EXAMPLE 72

4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0157] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,4-methylenedioxyaniline (53 mg, 0.386 mmol) similar to Example 58 and isolated as a light brown solid (37 mg, 53%). ^1H NMR (CDCl_3): 9.63 (dd, $J = 0.9, 1.8$ Hz, 1H), 8.73-8.68 (m, 2H), 7.43-7.39 (m, 1H), 6.93 (s, 1H), 6.88 (d, $J = 8.1$ Hz, 2H), 6.83 (dd, $J = 1.8, 8.1$ Hz, 1H), 6.77 (s, 1H), 6.06 (s, 2H).

EXAMPLE 73

4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

[0158] a. 4-Chloro-2-phenyl-6-(trifluoromethyl)pyrimidine: A mixture of 4-hydroxy-2-phenyl-6-(trifluoromethyl)pyrimidine (1.5 g, 6.24 mmol) and phosphorus oxychloride (25 ml) was refluxed for 3 h. The mixture was cooled to room temperature and then rotary evaporated to dryness. The residual was extracted with ethyl acetate (100 ml), washed with water (2 x 50 ml), and then with aqueous saturated sodium chloride (1 x 50 ml). The ethyl acetate solution was dried over anhydrous sodium sulfate and then rotary evaporated to dryness to give a pale white solid (1.6 g, 99%). ^1H NMR ($\text{DMSO}-d_6$): 8.39-8.36 (m, 2H), 8.26 (s, 1H), 7.66-7.57 (m, 3H).

[0159] b. 4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine: A mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), m-anisidine (33 μl , 0.290 mmol) and aqueous 2N HCl (150 μl) in water:ethanol (1:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and then basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (75 ml), washed with water (2 x 25

- 78 -

ml), and then with aqueous saturated sodium chloride (1 x 25 ml). The ethyl acetate solution was dried over anhydrous sodium sulfate and then rotary evaporated to dryness. The residual was purified by column chromatography and isolated as a light gray solid (15 mg, 23 %). ¹H NMR (CDCl₃): 8.47-8.44 (m, 2H), 7.51-7.46 (m, 3H), 7.34 (t, *J* = 8.1 Hz, 1H), 7.16 (s, 1H), 7.10 (s, 1H), 6.98 (dd, *J* = 1.5, 7.8 Hz, 1H), 6.89 (s, 1H), 6.81-6.77 (m, 1H), 3.85 (s, 3H).

EXAMPLE 74

4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

[0160] a. 4-Chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine: A mixture of 4-hydroxy-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (2.0 g, 8.26 mmol) and phosphorus oxychloride (30 ml) was refluxed for 3 h. The mixture was cooled to room temperature and then rotary evaporated to dryness. The residual was purified by column chromatography and isolated as a tan solid (1.8 g, 84%). ¹H NMR (DMSO-d₆): 9.54 (d, *J* = 1.5 Hz, 1H), 8.92 (dd, *J* = 1.5, 2.4 Hz, 1H), 8.90 (d, *J* = 2.4 Hz, 1H), 8.48 (s, 1H).

[0161] b. 4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine: A mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol), m-anisidine (32 μl, 0.288 mmol) and aqueous 2N HCl (150 μl) in water:ethanol (1:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and then basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (75 ml), washed with water (2 x 25 ml), and then with aqueous saturated sodium chloride (1 x 25 ml). The ethyl acetate solution was dried over anhydrous sodium sulfate and then rotary evaporated to dryness. The residual was purified by column chromatography and isolated as a yellow solid (38 mg, 57 %). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.2 Hz, 1H), 8.74 (dd, *J* = 1.5, 2.4 Hz, 1H), 8.71 (d, *J* = 2.4 Hz, 1H), 8.04 (s, 1H), 7.34 (t, *J* = 8.1 Hz,

- 79 -

1H), 7.07 (s, 1H), 7.01 (s, 1H), 6.94 (d, $J = 8.1$ Hz, 1H), 6.82 (dd, $J = 2.7, 8.4$ Hz, 1H), 3.84 (s, 3H).

EXAMPLE 75

4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

[0162] A mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 2,5-dimethoxyaniline (44 mg, 0.290 mmol) and aqueous 2N HCl (150 μ l) in water:ethanol (1:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and then basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water, water:ethanol (2:1) and dried under vacuo to give a gray solid (36 mg, 50%). ^1H NMR (CDCl_3): 8.51-8.48 (m, 2H), 8.14 (s, 1H), 7.51-7.48 (m, 4H), 6.89 (s, 1H), 6.88 (d, $J = 9.3$ Hz, 1H), 6.66 (dd, $J = 3.0, 8.7$ Hz, 1H), 3.90 (s, 3H), 3.87 (s, 3H).

EXAMPLE 76

4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

[0163] A mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol), 2,5-dimethoxyaniline and aqueous 2N HCl (150 μ l) in water:ethanol (1:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and then basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water, water:ethanol (2:1) and isolated as a green-yellow solid (46 mg, 64%). ^1H NMR (CDCl_3): 9.74 (d, $J = 1.5$ Hz, 1H), 8.79 (dd, $J = 1.8, 2.4$ Hz, 1H), 8.72 (d, $J = 2.4$ Hz, 1H), 7.80 (bs, 1H), 7.70 (s, 1H), 7.07 (s, 1H), 6.91 (d, $J = 9.3$ Hz, 1H), 6.72 (dd, $J = 3.0, 9.0$ Hz, 1H), 3.87 (s, 3H), 3.86 (s, 3H).

EXAMPLE 77

4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

- [0164] The title compound was prepared from a mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,4-dimethoxyaniline (44 mg, 0.290 mmol) similar to Example 75. The mixture was extracted with ethyl acetate (75 ml), washed with water (2 x 25 ml), washed with aqueous saturated NaCl (1 x 25 ml), and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness. The residual was purified by column chromatography and isolated as a yellow oil (35 mg, 48%). ¹H NMR (CDCl₃): 8.45-8.42 (m, 2H), 7.50-7.44 (m, 3H), 7.25 (s, 1H), 7.02 (s, 1H), 6.90 (s, 2H), 6.74 (s, 1H), 3.91 (s, 3H), 3.87 (s, 3H).

EXAMPLE 78

4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

- [0165] The title compound was prepared from a mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol), 3,4-dimethoxyaniline (44 mg, 0.288 mmol) similar to Example 76 and isolated as a yellow solid (52 mg, 72%). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.2 Hz, 1H), 8.78 (dd, *J* = 1.2, 2.4 Hz, 1H), 8.72 (d, *J* = 2.4 Hz, 1H), 7.36 (s, 1H), 6.96-6.90 (m, 3H), 6.88 (s, 1H), 3.94 (s, 3H), 3.91 (s, 3H).

EXAMPLE 79

4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

- [0166] The title compound was prepared from a mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-methoxy-2-methylaniline (40 mg, 0.290 mmol) similar to Example 77 and isolated as a tan oil (33 mg, 48%). ¹H NMR (CDCl₃): 8.46-8.42 (m, 2H), 7.50-7.46 (m,

- 81 -

3H), 7.22 (d, $J = 8.4$ Hz, 1H), 7.01 (d, $J = 2.4$ Hz, 1H), 6.92 (s, 1H), 6.80 (dd, $J = 2.6, 8.6$ Hz, 1H), 6.61 (s, 1H), 3.81 (s, 3H), 2.22 (s, 3H).

EXAMPLE 80

4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

[0167] The title compound was prepared from a mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol), 5-methoxy-2-methylaniline (40 mg, 0.288 mmol) similar to Example 76 and isolated as a yellow solid (40 mg, 58%). ^1H NMR (CDCl_3): 9.75 (d, $J = 1.2$ Hz, 1H), 8.79 (dd, $J = 1.5, 2.4$ Hz, 1H), 8.73 (d, $J = 2.4$ Hz, 1H), 7.28 (d, $J = 4.5$ Hz, 1H), 7.24 (s, 1H), 6.89 (d, $J = 2.7$ Hz, 1H), 6.86 (dd, $J = 2.7, 8.4$ Hz, 1H), 6.73 (s, 1H), 3.82 (s, 3H), 2.22 (s, 3H).

EXAMPLE 81

4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

[0168] A mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-chloro-5-methoxyaniline hydrochloride (56 mg, 0.290 mmol) in water:ethanol (1:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and then basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water, water:ethanol (2:1) and isolated as a brown solid (28 mg, 38%). ^1H NMR (CDCl_3): 8.50-8.47 (m, 2H), 7.99 (d, $J = 3.0$ Hz, 1H), 7.53-7.47 (m, 3H), 7.36 (d, $J = 9.0$ Hz, 1H), 7.29 (s, 1H), 6.91 (s, 1H), 6.71 (dd, $J = 3.0, 9.0$ Hz, 1H), 3.88 (s, 3H).

EXAMPLE 82

4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

[0169] The title compound was prepared from a mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol) and 2-chloro-

- 82 -

5-methoxyaniline hydrochloride (56 mg, 0.288 mmol) similar to Example 76 and isolated as a tan solid (41 mg, 56%). ¹H NMR (CDCl₃): 9.75 (d, *J* = 1.2 Hz, 1H), 8.79 (dd, *J* = 1.7, 2.6 Hz, 1H), 8.73 (d, *J* = 2.4 Hz, 1H), 7.70 (s, 1H), 7.52 (s, 1H), 7.40 (d, *J* = 9.3 Hz, 1H), 7.04 (s, 1H), 6.78 (dd, *J* = 2.9, 8.9 Hz, 1H), 3.89 (s, 3H).

EXAMPLE 83

4-(3,4-Methylenedioxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine

[0170] The title compound was prepared from a mixture of 4-chloro-2-phenyl-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,4-methylenedioxyaniline (40 mg, 0.290 mmol) similar to Example 77 and isolated as a tan solid (16 mg, 23%). ¹H NMR (CDCl₃): 8.44-8.40 (m, 2H), 7.50-7.45 (m, 3H), 7.02 (s, 1H), 6.91 (s, 1H), 6.85 (d, *J* = 8.1 Hz, 1H), 6.78 (dd, *J* = 2.0, 8.3 Hz, 1H), 6.71 (s, 1H), 6.03 (s, 2H).

EXAMPLE 84

4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine

[0171] The title compound was prepared from a mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol) and 3,4-methylenedioxyaniline (39 mg, 0.288 mmol) similar to Example 76 and isolated as a pink solid (21 mg, 30%). ¹H NMR (CDCl₃): 9.72 (d, *J* = 1.2 Hz, 1H), 8.76 (dd, *J* = 1.5, 2.7 Hz, 1H), 8.71 (d, *J* = 2.7 Hz, 1H), 7.60 (s, 1H), 6.88 (s, 1H), 6.87 (d, *J* = 8.1 Hz, 1H), 6.83 (s, 1H), 6.78 (dd, *J* = 1.8, 8.1 Hz, 1H), 6.05 (s, 2H).

EXAMPLE 85

4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0172] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-methoxy-5-phenoxyaniline (62 mg, 0.290 mmol) similar to Example 58 and isolated as a tan solid (46 mg, 54%). ¹H NMR (CDCl₃): 9.57 (dd, *J* = 1.1, 2.3 Hz, 1H), 8.67 (dd, *J* = 1.8, 4.8 Hz, 1H), 8.42-8.38 (m, 1H), 8.12 (s, 1H), 7.75 (s, 1H), 7.38-7.31 (m, 2H), 7.25-7.21 (m, 1H), 7.11 (t, *J* = 1.2 Hz, 1H), 7.09-7.03 (m, 2H), 6.91 (d, *J* = 8.7 Hz, 2H), 6.82 (dd, *J* = 2.7, 8.7 Hz, 1H), 3.91 (s, 3H).

EXAMPLE 86

4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0173] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-amino-4-methylbenzoate (48 mg, 0.290 mmol) similar to Example 58 and isolated as a yellow solid (2 mg, 3%). ¹H NMR (CDCl₃): 9.61 (s, 1H), 8.69-8.65 (m, 2H), 8.22 (s, 1H), 7.90 (dd, *J* = 1.8, 7.8 Hz, 1H), 7.40-7.35 (m, 2H), 7.14 (s, 1H), 6.59 (s, 1H), 3.91 (s, 3H), 2.35 (s, 3H).

EXAMPLE 87

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0174] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-methoxy-5-methylaniline (40 mg, 0.290 mmol) similar to Example 58 and isolated as a yellow solid (41 mg, 59%). ¹H NMR (CDCl₃): 9.66 (d, *J* = 1.8

- 84 -

Hz, 1H), 8.74-8.70 (m, 2H), 7.98 (s, 1H), 7.55 (s, 1H), 7.44-7.40 (m, 1H), 6.99-6.95 (m, 1H), 6.88 (s, 1H), 6.85 (s, 1H), 3.89 (s, 3H), 2.40 (s, 3H).

EXAMPLE 88

4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0175] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-fluoro-2-methylaniline (36 mg, 0.290 mmol) similar to Example 58 and isolated as a tan solid (4 mg, 6%). ¹H NMR (CDCl₃): 9.71 (dd, *J* = 0.80, 2.3 Hz, 1H), 8.74-8.71 (m, 1H), 8.68 (dd, *J* = 1.7, 5.0 Hz, 1H), 7.72 (s, 1H), 7.44-7.39 (m, 1H), 7.33-7.28 (m, 2H), 7.02-6.95 (m, 1H), 6.67 (s, 1H), 2.29 (s, 3H).

EXAMPLE 89

4-(2-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0176] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-amino-4-methoxybenzotrifluoride (55 mg, 0.290 mmol) similar to Example 58 and isolated as a yellow solid (56 mg, 70%). ¹H NMR (CDCl₃): 9.66 (dd, *J* = 0.6, 2.1 Hz, 1H), 8.95 (s, 1H), 8.75 (dd, *J* = 1.8, 4.8 Hz, 1H), 8.73-8.69 (m, 1H), 7.67 (s, 1H), 7.46-7.42 (m, 1H), 7.41-7.37 (m, 1H), 7.01 (d, *J* = 8.4 Hz, 1H), 6.93 (s, 1H), 3.99 (s, 3H).

EXAMPLE 90

4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0177] A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (500 mg, 1.93 mmol), 2-methyl-5-nitroaniline (441 mg, 2.90 mmol), and 2N HCl (1.5 ml) in water:ethanol (1:1, 100 ml) was refluxed for 48 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH

- 85 -

10-12. The resulting precipitate was filtered, washed with water and water:ethanol (2:1). The product was dried under vacuo and then washed with chloroform to give a yellow solid (136 mg, 19%). ¹H NMR (DMSO-d₆): 9.84 (s, 1H), 9.41 (s, 1H), 8.99 (s, 1H), 8.73 (d, *J* = 4.2 Hz, 1H), 8.57 (d, *J* = 7.8 Hz, 1H), 8.02 (dd, *J* = 1.8, 7.8, 1H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.58-7.54 (m, 1H), 7.39 (s, 1H), 2.45 (s, 3H).

EXAMPLE 91

4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0178] To a solution of 4-(2-methyl,5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (75 mg, 0.200 mmol) in anhydrous ethanol (10 ml) was added slowly anhydrous tin chloride (152 mg, 0.800 mmol). The mixture was refluxed for 1 h and then rotary evaporated to leave a brown oil. The brown oil was basified with 2N NaOH to give a final pH of 8. The resulting precipitate was filtered and washed with ethyl acetate. The aqueous phase and organic phase was combined, washed with water (1x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate was rotary evaporated to dryness to give a dark yellow solid (33 mg, 44%). ¹H NMR (CDCl₃): 9.69-9.68 (m, 1H), 8.73-8.67 (m, 2H), 7.53 (s, 1H), 7.43-7.39 (m, 1H), 7.11 (d, *J* = 8.4 Hz, 1H), 6.72 (s, 1H), 6.64 (d, *J* = 2.7 Hz, 1H), 6.61 (s, 1H), 3.01 (bs, 2H), 2.17 (s, 3H).

EXAMPLE 92

4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0179] A mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol), m-phenetidine (19 µl, 0.144 mmol), and 2N HCl (75 µl) in water:ethanol (2:1, 5 ml) was refluxed for 24 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water, water:ethanol (2:1) and

- 86 -

dried to give a yellow solid (22 mg, 63%). ^1H NMR (CDCl_3): 8.88-8.85 (m, 1H), 8.58 (d, $J = 8.1$ Hz, 1H), 7.93-7.88 (m, 1H), 7.61 (s, 1H), 7.48-7.44 (m, 1H), 7.35 (t, $J = 8.4$ Hz, 1H), 7.05 (s, 1H), 6.89-6.81 (m, 3H), 4.07 (q, $J = 7.1$ Hz, 2H), 1.45 (t, $J = 7.1$ Hz, 3H).

EXAMPLE 93

4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0180] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol) and 3-ethylaniline (18 μl , 0.144 mmol) similar to Example 92 and isolated as a yellow oil (12 mg, 36%). ^1H NMR (CDCl_3): 8.84-8.82 (m, 1H), 8.57-8.54 (m, 1H), 7.89-7.84 (m, 1H), 7.72 (s, 1H), 7.44-7.40 (m, 1H), 7.34 (d, $J = 8.7$ Hz, 1H), 7.14-7.11 (m, 3H), 7.01 (s, 1H), 2.69 (q, $J = 7.8$ Hz, 2H), 1.27 (t, $J = 7.8$ Hz, 3H).

EXAMPLE 94

4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0181] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol) and 5-methoxy-2-methylaniline (20 mg, 0.144 mmol) similar to Example 92 and isolated as a yellow solid (18 mg, 52%). ^1H NMR (CDCl_3): 8.85-8.82 (m, 1H), 8.59-8.56 (m, 1H), 7.90-7.84 (m, 1H), 7.45-7.40 (m, 2H), 7.24 (d, $J = 8.1$ Hz, 1H), 6.87 (s, 1H), 6.83 (dd, $J = 2.6, 8.3$ Hz, 1H), 6.69 (s, 1H), 3.81 (s, 3H), 2.20 (s, 3H).

- 87 -

EXAMPLE 95

4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0182] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol) and 2-chloro-5-methoxyaniline (23 mg, 0.144 mmol) similar to Example 92 and isolated as a white solid (11 mg, 48%). ¹H NMR (CDCl₃): 8.84-8.82 (m, 1H), 8.58-8.54 (m, 1H), 7.90-7.85 (m, 1H), 7.66 (s, 1H), 7.60 (s, 1H), 7.46-7.41 (m, 1H), 7.37 (d, *J* = 8.7 Hz, 1H), 7.01 (s, 1H), 6.75 (dd, *J* = 3.0, 8.7 Hz, 1H), 3.88 (s, 3H).

EXAMPLE 96

4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0183] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol) and 3-(methylmercapto)aniline (18 μl, 0.144 mmol) similar to Example 92 and isolated as a yellow solid (21 mg, 58%). ¹H NMR (CDCl₃): 8.87-8.85 (m, 1H), 8.58 (d, *J* = 7.8 Hz, 1H), 7.93-7.88 (m, 1H), 7.61 (s, 1H), 7.48-7.44 (m, 1H), 7.36 (t, *J* = 7.8 Hz, 1H), 7.25 (s, 1H), 7.17-7.14 (m, 1H), 7.10-7.06 (m, 1H), 7.01 (s, 1H), 2.52 (s, 3H).

EXAMPLE 97

4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride

[0184] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 3-aminophenol (32 mg, 0.290 mmol), and 2N HCl (150 μl) in water:ethanol (2:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature and the resulting crystals was filtered, washed with water, with water:ethanol (2:1) and dried to give a tan crystals (22 mg, 30%). ¹H NMR (DMSO-d₆): 10.17 (s, 1H), 9.36 (d, *J* = 1.5 Hz, 1H), 8.72-8.66 (m, 2H), 7.66-

- 88 -

7.62 (m, 1H), 7.20 (s, 1H), 7.10 (d, $J = 7.8$ Hz, 1H), 7.05 (d, $J = 7.8$ Hz, 1H), 7.04 (s, 1H), 7.01 (s, 1H), 6.45-6.41 (m, 1H).

EXAMPLE 98

4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride

[0185] The title compound was prepared from a mixture of 4-chloro-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.192 mmol) and 3-aminophenol (31 mg, 0.288 mmol) similar to Example 97 and isolated as a tan solid (28 mg,). ^1H NMR (DMSO- d_6): 10.26 (s, 1H), 9.62 (s, 1H), 9.51-9.50 (m, 1H), 8.87-8.83 (m, 2H), 7.30 (s, 1H), 7.23-7.21 (m, 3H), 6.57-6.54 (m, 1H).

EXAMPLE 99

4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride

[0186] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 3-aminophenol (32 mg, 0.290 mmol) similar to Example 97 and isolated as a yellow solid (37 mg, 52%). ^1H NMR (DMSO- d_6): 10.51 (s, 1H), 8.98 (d, $J = 6.0$ Hz, 2H), 8.54 (d, $J = 6.6$ Hz, 2H), 7.32-7.21 (m, 5H), 6.61-6.57 (m, 1H).

EXAMPLE 100

4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride

[0187] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 2,4-dimethoxyaniline (56 mg, 0.365 mmol) similar to Example 92 and isolated as a purple crystal (52 mg, 64%). ^1H NMR (CDCl_3): 8.84-8.81 (m, 1H), 8.48-8.44 (m, 1H), 7.86-7.80 (m, 1H), 7.60 (d, $J = 8.4$ Hz, 1H), 7.38-7.34 (m, 1H), 6.93 (s, 1H), 6.56-6.53

- 89 -

(m, 2H), 6.46 (s, 1H), 3.84 (d, $J = 0.6$ Hz, 3H), 3.83 (d, $J = 0.9$ Hz, 3H), 2.48 (s, 3H).

EXAMPLE 101

4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride

[0188] The title compound was prepared from a mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol) and 3,5-dimethoxyaniline (56 mg, 0.365 mmol) similar to Example 92 and isolated as a tan solid (21 mg, 26%). ^1H NMR (CDCl_3): 8.83-8.81 (m, 1H), 8.49-8.46 (m, 1H), 7.86-7.80 (m, 1H), 7.39-7.35 (m, 1H), 6.99 (s, 1H), 6.70 (s, 1H), 6.53 (d, $J = 2.4$ Hz, 2H), 6.30 (t, $J = 2.3$ Hz, 1H), 3.82 (s, 6H), 2.51 (s, 3H).

EXAMPLE 102

4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0189] A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 2,5-diethoxyaniline (52 mg, 0.290 mmol) and 2N HCl (150 μl) in water:ethanol (3:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered and isolated as a yellow solid (14 mg, 18%). ^1H NMR (CDCl_3): 9.67 (s, 1H), 8.74-8.71 (m, 2H), 7.93 (s, 1H), 7.64 (s, 1H), 7.43-7.39 (m, 1H), 6.91 (s, 1H), 6.86 (d, $J = 9.3$ Hz, 1H), 6.65 (dd, $J = 2.7, 8.7$ Hz, 1H), 4.13-4.04 (m, 4H), 1.47-1.42 (m, 6H).

EXAMPLE 103

4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0190] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-amino-4-methoxybenzoic acid (48 mg, 0.290 mmol) similar to Example 92 and isolated

- 90 -

as a white solid (50 mg, 64%). ¹H NMR (DMSO-d₆): 9.78 (s, 1H), 9.49-9.48 (m, 1H), 9.08 (s, 1H), 8.74 (dd, *J* = 1.5, 4.8 Hz, 1H), 8.70-8.67 (m, 1H), 7.79 (dd, *J* = 1.5, 4.8 Hz, 1H), 7.56-7.52 (m, 2H), 7.23 (d, *J* = 9.0 Hz, 1H), 3.97 (s, 3H).

EXAMPLE 104

4-(3-Methoxybenzylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0191] A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-methoxybenzylamine (37 μ l, 0.290 mmol) in 10 ml of ethanol was stirred at room temperature for 48 h. The mixture was rotary evaporated to dryness and purified by preparative TLC with hexane:ethyl acetate (3:1) to give a yellow solid (14 mg, 20%). ¹H NMR (CDCl₃): 9.60-9.59 (m, 1H), 8.70-8.66 (m, 2H), 7.40-7.32 (m, 1H), 7.30 (d, *J* = 7.8 Hz, 1H), 6.95 (d, *J* = 7.8 Hz, 1H), 6.91 (s, 1H), 6.85 (dd, *J* = 2.7, 8.4 Hz, 1H), 6.62 (s, 1H), 5.75 (bs, 1H), 4.73 (s, 2H), 3.79 (s, 3H).

EXAMPLE 105

4-(5-Carboxyl-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0192] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-amino-4-hydroxybenzoic acid (44 mg, 0.290 mmol) similar to Example 92 and isolated as a brown solid (25 mg, 34%). ¹H NMR (DMSO-d₆): 9.73 (s, 1H), 9.49 (d, *J* = 1.5 Hz, 1H), 9.07 (s, 1H), 8.74 (dd, *J* = 1.5, 4.5 Hz, 1H), 8.70-8.66 (m, 1H), 7.65 (dd, *J* = 2.4, 8.7 Hz, 1H), 7.56-7.52 (m, 2H), 7.03 (d, *J* = 8.4 Hz, 1H).

EXAMPLE 106

4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0193] a. 2-Chloro-5-hydroxyaniline: To a solution of 4-chloro-3-nitrophenol (1 g, 5.76 mmol) in anhydrous ethanol (20 ml) was slowly added anhydrous tin chloride (4.4 g, 23.2 mmol) at room temperature. The mixture was refluxed for 1 h. The mixture was cooled to room temperature and then rotary evaporated to get an oily liquid. The liquid was basified with aqueous 2N NaOH (20 ml) to a final pH of 8. The resulting precipitate was filtered and washed with water. The aqueous solution was extracted with ethyl acetate (75 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness to give a pale white solid (584 mg, 71%). ¹H NMR (CDCl₃): 7.07 (d, *J* = 8.4 Hz, 1H), 6.27 (d, *J* = 3.0 Hz, 1H), 6.18 (dd, *J* = 3.0, 9.0 Hz, 1H), 4.64 (s, 1H), 4.02 (bs, 2H).

[0194] b. 4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine: A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 2-chloro-5-hydroxyaniline (42 mg, 0.290 mmol) and 2N HCl (150 μl) in water:ethanol (2:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature, neutralized with aqueous 2N NaOH to a pH of 6-7 and isolated as a yellow solid (6 mg, 8%). ¹H NMR (DMSO-*d*₆): 9.95 (s, 1H), 9.85 (s, 1H), 9.38-9.37 (m, 1H), 8.74-8.72 (m, 1H), 8.55-8.52 (m, 1H), 7.59-7.54 (m, 1H), 7.37 (dd, *J* = 0.6, 8.7 Hz, 1H), 7.38 (s, 1H), 7.24 (s, 1H), 6.71 (dd, *J* = 2.1, 8.7 Hz, 1H).

EXAMPLE 107

4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0195] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-chloro-5-hydroxyaniline (42 mg, 0.290 mmol) similar to Example 108 and isolated as a brown solid (3 mg, 4%). ¹H NMR (CDCl₃): 8.68 (dd, *J* = 1.8, 4.5 Hz, 2H),

- 92 -

8.13 (dd, $J = 1.8, 4.2$ Hz, 2H), 7.99 (d, $J = 2.7$ Hz, 1H), 7.34 (s, 1H), 7.33 (d, $J = 8.7$ Hz, 1H), 6.98 (s, 1H), 6.73 (dd, $J = 2.7, 8.7$ Hz, 1H).

EXAMPLE 108

4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0196] A mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (50 mg, 0.243 mmol), 2,6-dimethylaniline (45 μ l, 0.365 mmol) and 2N HCl (150 μ l) in water:ethanol (2:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (75 ml), which was washed with water (2x20 ml), aqueous saturated NaCl (1x20 ml), and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness and the residue was purified by column chromatography to give a yellow oil (58 mg, 82%). ^1H NMR (CDCl_3): 8.81-8.78 (m, 1H), 8.47 (d, $J = 8.1$ Hz, 1H), 7.83-7.77 (m, 1H), 7.36-7.30 (m, 1H), 7.21-7.13 (m, 3H), 6.91 (s, 1H), 5.77 (s, 1H), 2.39 (s, 3H), 2.23 (s, 6H).

EXAMPLE 109

4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine

[0197] A mixture 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol), m-anisidine (16 μ l, 0.140 mmol) and 2N HCl (75 μ l) in water:ethanol (2:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was filtered, washed with water and aqueous ethanol to give a yellow solid (20 mg, 60%). ^1H NMR (CDCl_3): 8.87-8.85 (m, 1H), 8.69-8.61 (m, 3H), 7.95 (s, 1H), 7.92-7.84 (m, 2H), 7.44-7.36 (m, 2H), 7.33 (d, $J = 8.1$ Hz, 1H), 7.30 (s, 1H), 7.05 (t, $J = 2.1$ Hz, 1H), 6.98 (dd, $J = 2.1, 8.1$ Hz, 1H), 6.79-6.75 (m, 1H), 3.86 (s, 3H).

EXAMPLE 110

4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine

- [0198] The title compound was prepared from a mixture 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol) and 2,5-dimethoxyaniline (21 mg, 0.140 mmol) similar to Example 111 and isolated as a yellow oil (2 mg, 6%). ¹H NMR (CDCl₃): 8.86-8.84 (m, 1H), 8.70-8.64 (m, 3H), 8.11 (s, 1H), 7.92 (s, 1H), 7.91-7.85 (m, 2H), 7.65 (s, 1H), 7.43-7.36 (m, 2H), 6.87 (d, *J* = 9.0 Hz, 1H), 6.61 (dd, *J* = 3.0, 9.0 Hz, 1H), 3.89 (s, 3H), 3.86 (s, 3H).

EXAMPLE 111

4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine

- [0199] The title compound was prepared from a mixture 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol) and 5-methoxy-2-methylaniline (19 mg, 0.140 mmol) similar to Example 111 and isolated as a white solid (21 mg, 61%). ¹H NMR (CDCl₃): 8.88-8.86 (m, 1H), 8.71-8.60 (m, 3H), 7.92-7.83 (m, 2H), 7.62 (s, 1H), 7.44-7.34 (m, 2H), 7.22 (d, *J* = 8.4 Hz, 1H), 7.08 (d, *J* = 2.7 Hz, 1H), 7.06 (s, 1H), 6.78 (dd, *J* = 2.4, 8.4 Hz, 1H), 3.82 (s, 3H), 2.52 (s, 3H).

EXAMPLE 112

4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine

- [0200] The title compound was prepared from a mixture 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol) and 2-methoxy-5-methylaniline (19 mg, 0.140 mmol) similar to Example 111 and isolated as a yellow solid (12 mg, 35%). ¹H NMR (CDCl₃): 8.89-8.87 (m, 1H), 8.69-8.64 (m, 3H), 7.92-7.84 (m, 3H), 7.89 (s, 1H), 7.55 (s, 1H), 7.44-7.36 (m, 2H), 6.92 (dd, *J* = 1.8, 8.1 Hz, 1H), 6.85 (d, *J* = 8.1 Hz, 1H), 3.86 (s, 3H), 2.39 (s, 3H).

EXAMPLE 113

4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine

- [0201] The title compound was prepared from a mixture of 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol) and 2-chloro-5-methoxyaniline (27 mg, 0.140 mmol) similar to Example 111 and isolated as a white solid (20 mg, 55%). ¹H NMR (CDCl₃): 8.87-8.82 (m, 1H), 8.72-8.64 (m, 3H), 8.08 (d, *J* = 2.7 Hz, 1H), 7.92 (s, 1H), 7.90-7.86 (m, 2H), 7.45-7.38 (m, 3H), 7.33 (d, *J* = 8.7 Hz, 1H), 6.67 (dd, *J* = 3.0, 9.0 Hz, 1H), 3.91 (s, 3H).

EXAMPLE 114

4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine

- [0202] The title compound was prepared from a mixture 4-chloro-2,6-di(2-pyridinyl)pyrimidine (25 mg, 0.093 mmol) and 2,5-dimethylaniline (17 μl, 0.140 mmol) similar to Example 111 and isolated as a yellow oil (7 mg, 21%). ¹H NMR (CDCl₃): 8.88-8.86 (m, 1H), 8.71-8.68 (m, 1H), 8.64-8.58 (m, 2H), 7.92-7.82 (m, 2H), 7.50 (s, 1H), 7.43-7.32 (m, 2H), 7.23 (s, 1H), 7.21 (d, *J* = 7.5 Hz, 1H), 7.09 (s, 1H), 7.04 (dd, *J* = 1.8, 7.8 Hz, 1H), 2.36 (s, 3H), 2.27 (s, 3H).

EXAMPLE 115

4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

- [0203] A mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 2,5-dimethylaniline (36 μl, 0.290 mmol) and 2N HCl (150 μl) in water:ethanol (2:1, 10 ml) was refluxed for 24 h. The mixture was cooled to room temperature. The resulting precipitate was filtered, washed with water and aqueous ethanol to give a white solid (28 mg, 42%). ¹H NMR (CDCl₃): 8.77 (dd, *J* = 1.5, 4.5 Hz, 2H), 8.27 (dd, *J* = 1.5, 4.5 Hz, 2H), 7.24 (d,

- 95 -

$J = 7.8$ Hz, 1H), 7.18 (s, 1H), 7.11 (dd, $J = 1.8, 7.8$ Hz, 1H), 6.99 (s, 1H), 6.62 (s, 1H), 2.38 (s, 3H), 2.25 (s, 3H).

EXAMPLE 116

4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0204] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-methoxy-5-methylaniline (40 mg, 0.290 mmol) similar to Example 115 and isolated as a white solid (48 mg, 69%). ^1H NMR (CDCl_3): 8.79 (dd, $J = 1.8, 4.5$ Hz, 2H), 8.30 (dd, $J = 1.8, 4.5$ Hz, 2H), 7.97 (s, 1H), 7.46 (s, 1H), 6.99 (dd, $J = 1.5, 8.1$ Hz, 1H), 6.94 (s, 1H), 6.89 (d, $J = 8.4$ Hz, 1H), 3.90 (s, 3H), 2.41 (s, 3H).

EXAMPLE 117

4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0205] A mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol), 5-hydroxy-2-methylaniline (36 mg, 0.290 mmol) and 2N HCl (150 μl) in water:ethanol (2:1, 10ml) was refluxed for 24 h. The mixture was cooled to room temperature and neutralized with aqueous 2N NaOH to pH 6-7. The mixture was extracted with ethyl acetate (75 ml), washed with water (2x20 ml), aqueous saturated NaCl (1x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness and the residue was purified by column chromatography to give a yellow solid (18 mg, 27%). ^1H NMR (Acetone- d_6): 9.53-9.51 (m, 1H), 8.85 (s, 1H), 8.71 (dd, $J = 1.8, 4.8$ Hz, 1H), 8.67-8.63 (m, 1H), 8.49 (s, 1H), 7.53-7.48 (m, 1H), 7.24 (s, 1H), 7.17 (d, $J = 8.1$ Hz, 1H), 6.98 (s, 1H), 6.75 (dd, $J = 2.4, 8.4$ Hz, 1H), 2.23 (s, 3H).

EXAMPLE 118

4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0206] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-hydroxy-2-methylaniline (36 mg, 0.290 mmol) similar to Example 117 and isolated as a tan solid (19 mg, 28%). ¹H NMR (Acetone-d₆): 8.91 (s, 1H), 8.75 (dd, *J* = 1.2, 4.8 Hz, 2H), 8.55 (s, 1H), 8.22 (dd, *J* = 1.8, 4.8 Hz, 2H), 7.23 (s, 1H), 7.18 (d, *J* = 8.1 Hz, 1H), 7.02 (s, 1H), 6.76 (dd, *J* = 3.0, 8.7 Hz, 1H), 2.23 (s, 3H).

EXAMPLE 119

4-(5-Methoxy-2-piperidino-anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0207] 1-(4-Methoxy-2-aminophenyl)piperidine: To a solution of 1-(4-methoxy-2-nitrophenyl)piperidine (238 mg, 1.01 mmol) in anhydrous ethanol (20 ml) was slowly added anhydrous tin chloride (772 mg, 4.07 mmol) at room temperature. The mixture was refluxed for 1 h. The mixture was cooled to room temperature and then rotary evaporated to get an oily liquid. The liquid was basified with aqueous 2N NaOH (20 ml) to a final pH of 10. The resulting precipitate was filtered and washed with water. The aqueous solution was extracted with ethyl acetate (75 ml), washed with water (2x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to give a dark brown liquid (153 mg, 74%). ¹H NMR (CDCl₃): 6.90 (d, *J* = 8.7 Hz, 1H), 6.29 (d, *J* = 2.7 Hz, 1H), 6.26 (dd, *J* = 2.7, 8.1 Hz, 1H), 4.04 (bs, 2H), 3.69 (s, 3H), 2.75 (s, 4H), 1.70-1.63 (m, 4H), 1.54 (s, 2H).

[0208] b. 4-(5-Methoxy-2-piperidino-anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine: The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 1-(4-methoxy-2-aminophenyl)piperidine (60 mg, 0.290 mmol) similar to Example 117 and isolated as a green solid (1 mg, 1%). ¹H NMR

- 97 -

(CDCl₃): 9.71-9.69 (m, 1H), 8.78-8.74 (m, 2H), 8.70 (s, 1H), 8.18 (s, 1H), 7.46-7.41 (m, 1H), 7.16 (d, *J* = 8.7 Hz, 1H), 6.90 (s, 1H), 6.66 (dd, *J* = 2.7, 8.4 Hz, 1H), 3.90 (s, 3H), 2.83-2.79 (m, 4H), 1.81-1.74 (m, 4H), 1.62 (s, 2H).

EXAMPLE 120

4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0209] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-cyano-5-methylaniline (38 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (8 mg, 12%). ¹H NMR (CDCl₃): 9.62-9.61 (m, 1H), 8.75-8.69 (m, 2H), 8.08 (s, 1H), 7.62 (s, 1H), 7.60 (d, *J* = 8.1 Hz, 1H), 7.46-7.41 (m, 1H), 7.12 (dd, *J* = 0.9, 8.1 Hz, 1H), 7.00 (s, 1H), 2.52 (s, 3H).

EXAMPLE 121

4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0210] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,5-dimethylaniline (36 μl, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (26 mg, 39%). ¹H NMR (CDCl₃): 9.66-9.64 (m, 1H), 8.73 (s, 1H), 8.72-8.69 (m, 1H), 7.44-7.40 (m, 1H), 7.30 (s, 1H), 7.04 (s, 2H), 6.93 (s, 1H), 6.90 (s, 1H), 2.38 (s, 3H), 2.37 (s, 3H).

EXAMPLE 122

4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine
hydrochloride

[0211] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3,5-dimethoxyaniline (44 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (29 mg, 38%). ¹H NMR (DMSO-d₆): 10.32 (s, 1H), 9.48-9.46

- 98 -

(m, 1H), 8.76-8.74 (m, 1H), 8.62-8.58 (m, 1H), 7.63-7.58 (m, 1H), 7.14 (s, 1H), 7.06 (s, 2H), 6.32 (s, 1H), 3.80 (s, 6H).

EXAMPLE 123

4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0212] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-chloro-5-methylaniline (41 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (15 mg, 21%). ¹H NMR (CDCl₃): 9.62-9.61 (m, 1H), 8.73-8.69 (m, 2H), 7.89 (s, 1H), 7.44-7.40 (m, 1H), 7.37 (d, *J* = 8.1 Hz, 1H), 7.30 (s, 1H), 7.00 (dd, *J* = 0.9, 8.4 Hz, 1H), 6.88 (s, 1H), 2.42 (s, 3H).

EXAMPLE 124

4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0213] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-methoxy-5-trifluoromethylaniline (55 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (27 mg, 34%). ¹H NMR (CDCl₃): 9.65-9.64 (m, 1H), 8.75-8.70 (m, 2H), 7.46-7.41 (m, 2H), 7.36 (s, 1H), 7.31 (s, 1H), 7.00 (s, 1H), 6.93 (s, 1H), 3.91 (s, 3H).

EXAMPLE 125

4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine

[0214] 4-Hydroxy-6-Trifluoromethyl-2-(6-trifluoromethyl-3-pyridinyl)-pyrimidine: A mixture of 6-(trifluoromethyl)pyridine-3-amidine (1 g, 5.26 mmol) and ethyl 4,4,4-trifluoro-2-butyrate (1.1 ml, 7.89 mmol) in ethanol (100 ml) was stirred at 0 °C for 20 min. The temperature of the mixture was

raised to 60 °C and was let to stir at this temperature for 30 min. Then aqueous KOH (0.86 g in 20 ml water) was added dropwise. The mixture was refluxed for 4 h. The mixture was cooled to room temperature, rotary evaporated and isolated as a white solid (1.5 g, 94%). ¹H NMR (CDCl₃): 9.72 (d, *J* = 2.4 Hz, 1H), 8.87 (dd, *J* = 2.1, 8.7 Hz, 1H), 7.93 (d, *J* = 8.1 Hz, 1H), 7.00 (s, 1H).

[0215] 4-Chloro-2-[6-(trifluoromethyl)-3-pyridinyl]-6-

(trifluoromethyl)pyrimidine: A mixture of 4-hydroxy-6-trifluoromethyl-2-(6-trifluoromethyl-3-pyridinyl)-pyrimidine (145 mg, 0.469 mmol) and phosphorus oxychloride (10 ml) was refluxed for 3 h. The mixture was cooled to room temperature and rotary evaporated to leave an oily residue. The residue was extracted with ethyl acetate (50 ml), washed with water (2x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was concentrated in vacuo to give a crude solid (80 mg).

[0216] 4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine: A mixture of crude 4-chloro-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine (50 mg, 0.152 mmol), 2-chloro-5-methoxyaniline (50 mg, 0.229 mmol) and 2N HCl (150 µl) in water:ethanol (2:1, 10 ml) was refluxed for 120 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (75 ml), washed with water (2x20 ml), with aqueous saturated NaCl (1x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness and the residue was purified by preparative TLC to give a white solid (3 mg, 4%). ¹H NMR (CDCl₃): 9.72 (s, 1H), 8.90 (dd, *J* = 3.0, 7.8 Hz, 1H), 7.81 (d, *J* = 8.1 Hz, 1H), 7.75 (s, 1H), 7.40 (d, *J* = 9.3 Hz, 1H), 7.36 (s, 1H), 6.99 (s, 1H), 6.75 (dd, *J* = 3.0, 8.7 Hz, 1H), 3.86 (s, 3H).

EXAMPLE 126

4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0217] a. 5-Bromo-2-methoxyaniline: To a solution of 4-bromo-2-nitroanisole (500 mg, 2.15 mmol) in anhydrous ethanol (15 ml) was added slowly anhydrous tin chloride (1.6 g, 8.62 mmol). The mixture was refluxed for 1h, then rotary evaporated to leave a brown oil which was neutralized with aqueous 2N NaOH (5 ml). The resulting solid was filtered and washed with ethyl acetate. The aqueous and ethyl acetate solutions were combined and washed with water. The ethyl acetate solution was rotary evaporated to dryness and the residue was purified column chromatography to give a white solid (46 mg, 11%). ¹H NMR (CDCl₃): 6.79 (dd, *J* = 2.1, 9.0 Hz, 1H), 6.80 (d, *J* = 1.8 Hz, 1H), 6.61 (d, *J* = 9.3 Hz, 1H), 3.84 (s, 2H), 3.80 (s, 3H).
- [0218] b. 4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine: The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-bromo-2-methoxyaniline (39 mg, 0.193 mmol) similar to Example 117 and isolated as a tan solid (26 mg, 32%). ¹H NMR (CDCl₃): 9.67-9.66 (m, 1H), 8.77-8.71 (m, 3H), 7.55 (s, 1H), 7.48-7.43 (m, 1H), 7.24 (dd, *J* = 2.4, 8.7 Hz, 1H), 6.92 (s, 1H), 6.83 (d, *J* = 9.0 Hz, 1H), 3.93 (s, 3H).

EXAMPLE 127

4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0219] a. 2-Bromo-5-methoxyaniline: The title compound was prepared from 4-bromo-3-nitroanisole (490 mg, 2.11 mmol), anhydrous ethanol (15 ml) and anhydrous tin chloride (1.6 g, 8.44 mmol) similar to Example 126a and isolated as a white solid (42 mg, 10%). ¹H NMR (CDCl₃): 7.26 (d, *J* = 8.7 Hz, 1H), 6.31 (d, *J* = 3.0 Hz, 1H), 6.22 (dd, *J* = 2.7, 9.0 Hz, 1H), 4.06 (s, 2H), 3.73 (s, 3H).

- [0220] b. 4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine: The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-bromo-5-methoxyaniline (39 mg, 0.193 mmol) similar to Example 117 and isolated as a yellow solid (2 mg, 2%). ¹H NMR (CDCl₃): 9.66 (s, 1H), 8.74-8.71 (m, 2H), 7.80 (s, 1H), 7.53 (d, *J* = 9.0 Hz, 1H), 7.48 (s, 1H), 7.45-7.41 (m, 1H), 6.93 (s, 1H), 6.69 (dd, *J* = 3.0, 8.7 Hz, 1H), 3.87 (s, 3H).

EXAMPLE 128

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0221] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 3-methyl-5-(trifluoromethyl)aniline (51 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (60 mg, 78%). ¹H NMR (DMSO-d₆): 10.60 (s, 1H), 9.45-9.44 (m, 1H), 8.77-8.75 (m, 1H), 8.60-8.56 (m, 1H), 8.30 (s, 1H), 7.68 (s, 1H), 7.62-7.58 (m, 1H), 7.32 (s, 1H), 7.16 (s, 1H), 2.45 (s, 3H).

EXAMPLE 129

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

- [0222] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-chloro-2-hydroxyaniline (42 mg, 0.290 mmol) similar to Example 117 and isolated as a tan solid (38 mg, 54%). ¹H NMR (DMSO-d₆): 10.44 (s, 1H), 9.74 (s, 1H), 9.42 (s, 1H), 8.75-8.74 (m, 1H), 8.58-8.54 (m, 1H), 8.29 (s, 1H), 7.48 (s, 1H), 7.61-7.57 (m, 1H), 7.08 (dd, *J* = 2.1, 8.7 Hz, 1H), 6.96 (d, *J* = 9.0 Hz, 1H).

- 102 -

EXAMPLE 130

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0223] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-chloro-2,5-dimethoxyaniline (54 mg, 0.290 mmol) similar to Example 117 and isolated as a brown solid (52 mg, 63%). ¹H NMR (CDCl₃): 9.66-9.64 (m, 1H), 8.76-8.71 (m, 2H), 8.35 (s, 1H), 7.54 (s, 1H), 7.45-7.41 (m, 1H), 7.01 (s, 1H), 6.92 (s, 1H).

EXAMPLE 131

4-(6-Chloro-3-methoxyanilino)-2-morpholino-6-(methyl)pyrimidine

[0224] The title compound was prepared from a mixture of 4-(4-chloro-6-methyl-pyrimidin-2-yl)-morpholine (23 mg, 0.108 mmol) and 6-chloro-m-anisidine (31 mg, 0.161 mmol) similar to Example 117 and isolated as a white solid (11 mg, 31%). ¹H NMR (CDCl₃): 8.27 (d, *J* = 3.0 Hz, 1H), 7.91 (s, 1H), 7.27 (d, *J* = 9.0 Hz, 1H), 6.97 (s, 1H), 6.54 (dd, *J* = 3.0, 8.7 Hz, 1H), 3.80 (s, 3H), 3.77 (s, 8H), 2.14 (s, 3H).

EXAMPLE 132

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0225] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2,4-dichloro-5-methoxyaniline (44 mg, 0.232 mmol) similar to Example 117, and isolated as a brown solid (9 mg, 11%). ¹H NMR (CDCl₃): 9.64-9.63 (m, 1H), 8.76-8.70 (m, 2H), 8.12 (s, 1H), 7.49 (s, 1H), 7.46-7.41 (m, 2H), 6.95 (s, 1H), 4.00 (s, 3H).

- 103 -

EXAMPLE 133

4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0226] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-aminoindole (38 mg, 0.290 mmol) similar to Example 117 and isolated as a tan solid (6 mg, 9%). ¹H NMR (CDCl₃): 9.75-9.74 (m, 1H), 8.77-8.73 (m, 1H), 8.69 (dd, *J* = 1.8, 5.1 Hz, 1H), 8.57 (s, 1H), 8.08 (s, 1H), 7.44-7.38 (m, 2H), 7.31-7.23 (m, 3H), 6.85 (s, 1H), 6.50-6.48 (m, 1H).

EXAMPLE 134

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0227] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-amino-4,5-dimethoxyacetophenone (56 mg, 0.290 mmol) similar to Example 117 and isolated as a tan solid (24 mg, 30%). ¹H NMR (CDCl₃): 12.49 (s, 1H), 9.65-9.64 (m, 1H), 8.86 (s, 1H), 8.74-8.70 (m, 2H), 7.45-7.41 (m, 1H), 7.35 (s, 1H), 6.99 (s, 1H), 4.12 (s, 3H), 3.95 (s, 3H), 2.68 (s, 3H).

EXAMPLE 135

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0228] A mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (1 g, 3.90 mmol), 2-methyl-5-nitroaniline (890 mg, 5.90 mmol), and 2N HCl (3 ml) in water:ethanol (1:1, 200 ml) was refluxed for 48 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The mixture was extracted with ethyl acetate (200 ml), washed with water (2x100 ml), with aqueous saturated NaCl (1x 100 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness to give a tan solid (377 mg, 30%). ¹H NMR (CDCl₃): 8.90 (s, 1H),

- 104 -

8.80 (dd, $J = 1.8, 4.8$ Hz, 2H), 8.30 (dd, $J = 1.5, 4.5$ Hz, 2H), 8.09 (dd, $J = 2.4, 8.1$ Hz, 1H), 7.51 (d, $J = 8.4$ Hz, 1H), 6.99 (s, 1H), 6.87 (s, 1H), 2.47 (s, 3H).

EXAMPLE 136

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0229] To a solution of 4-(2-methyl,5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (377 mg, 1.00 mmol) in anhydrous ethanol (20 ml) was added slowly anhydrous tin chloride (758 mg, 4.00 mmol). The mixture was refluxed for 1 h and then rotary evaporated to leave a brown oil. The brown oil was basified with aqueous 2N NaOH to give a final pH of 8. The resulting precipitate was filtered and washed with ethyl acetate. The aqueous phase and organic phase was combined and washed with water (1x20 ml) and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness to give a brown solid (212 mg, 57%). ^1H NMR (CDCl_3): 8.76 (dd, $J = 1.8, 4.8$ Hz, 2H), 8.26 (dd, $J = 1.8, 4.8$ Hz, 2H), 7.11 (d, $J = 7.8$ Hz, 1H), 7.12 (s, 1H), 6.69 (d, $J = 2.4$ Hz, 1H), 6.66 (s, 1H), 6.63 (dd, $J = 2.4, 8.1$ Hz, 1H), 3.72 (bs, 2H), 2.16 (s, 3H).

EXAMPLE 137

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0230] A mixture of 4-(5-amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride (25 mg, 0.067 mmol), benzoic acid (10 mg, 0.080 mmol), and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (15.4 mg, 0.080 mmol) in dichloromethane (5 ml) was stirred at room temperature for 24 h. The mixture was rotary evaporated to dryness and the residue was purified by column chromatography to give a white solid (14 mg, 47%). ^1H NMR (CDCl_3): 9.66 (s, 1H), 8.75-8.71 (m, 1H), 8.67 (bs, 1H), 8.40 (s, 1H), 8.08 (dd, $J = 1.2, 8.4$ Hz, 1H), 7.93 (d, $J = 1.8$ Hz, 1H), 7.90-7.86

- 105 -

(m, 2H), 7.81 (s, 1H), 7.55-7.38 (m, 4H), 7.25 (d, $J = 8.4$ Hz, 1H), 6.71 (s, 1H), 2.23 (s, 3H).

EXAMPLE 138

4-(2,5-Diethoxy-4-morpholinoanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0231] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2,5-diethoxy-4-morpholinoaniline dihydrochloride (98 mg, 0.290 mmol) similar to Example 117 and isolated as a brown oil (3 mg, 3%). ^1H NMR (CDCl_3): 9.64-9.63 (m, 1H), 8.74-8.69 (m, 2H), 7.42-7.38 (m, 1H), 7.33 (s, 1H), 6.83 (s, 1H), 6.60 (s, 1H), 4.13 (q, $J = 7.2$ Hz, 4H), 3.92-3.89 (m, 4H), 3.14-3.11 (m, 4H), 1.26 (t, $J = 7.2$ Hz, 6H).

EXAMPLE 139

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0232] A mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (20 mg, 0.058 mmol) and p-toluoyl chloride in pyridine (5 ml) was stirred at room temperature for 15 h. Then water was added to the mixture. The resulting precipitate was filtered, washed with excess water and dried to give a tan solid (26 mg, 94%). ^1H NMR (CDCl_3): 8.77 (dd, $J = 1.8, 4.5$ Hz, 2H), 8.29 (dd, $J = 1.8, 4.5$ Hz, 2H), 8.00 (s, 1H), 7.83 (s, 1H), 7.78 (d, $J = 8.1$ Hz, 2H), 7.38 (dd, $J = 2.1, 8.4$ Hz, 1H), 7.34 (s, 1H), 7.31 (d, $J = 7.5$ Hz, 2H), 7.01 (s, 1H), 6.73 (s, 1H), 2.44 (s, 3H), 2.29 (s, 3H).

- 106 -

EXAMPLE 140

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide

- [0233] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (20 mg, 0.058 mmol) and nicotinoyl chloride hydrochloride (10 mg, 0.058 mmol) similar to Example 139 and isolated as a tan solid (13 mg, 47%). ¹H NMR (DMSO-d₆): 10.50 (s, 1H), 9.85 (s, 1H), 9.12 (d, *J* = 1.5 Hz, 1H), 8.77 (dd, *J* = 1.5, 4.5 Hz, 1H), 8.72 (d, *J* = 5.7 Hz, 2H), 8.34-8.28 (m, 1H), 8.21 (bs, 1H), 8.16 (d, *J* = 6.0 Hz, 2H), 7.60-7.53 (m, 2H), 7.34 (d, *J* = 8.1 Hz, 1H), 7.14 (s, 1H), 2.25 (s, 3H).

EXAMPLE 141

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide

- [0234] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (20 mg, 0.058 mmol) and 6-chloronicotinyl chloride (10 mg, 0.058 mmol) similar to Example 139 and isolated as a white solid (14 mg, 50%). ¹H NMR (CDCl₃): 8.88-8.87 (m, 1H), 8.77 (dd, *J* = 1.8, 4.5 Hz, 2H), 8.29 (dd, *J* = 1.8, 4.5 Hz, 2H), 8.19 (dd, *J* = 2.4, 8.4 Hz, 1H), 8.00 (s, 1H), 7.87 (s, 1H), 7.49 (dd, *J* = 0.9, 8.4 Hz, 1H), 7.36 (s, 1H), 7.35 (s, 1H), 7.01 (s, 1H), 6.76 (s, 1H), 2.32 (s, 3H).

EXAMPLE 142

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide

- [0235] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (10 mg, 0.029

- 107 -

mmol) and 4-morpholinobenzoic acid (7 mg, 0.035 mmol) similar to Example 136 and isolated as a white solid (4 mg, 26%). ¹H NMR (CDCl₃): 8.76 (dd, *J* = 1.8, 4.2 Hz, 2H), 8.29 (dd, *J* = 1.8, 4.2 Hz, 2H), 7.99 (s, 1H), 7.95 (s, 1H), 7.82 (d, *J* = 9.0 Hz, 2H), 7.37 (dd, *J* = 1.8, 8.1 Hz, 1H), 7.31 (d, *J* = 8.7 Hz, 1H), 7.07 (s, 1H), 6.93 (d, *J* = 9.0 Hz, 2H), 6.73 (s, 1H), 3.89-3.86 (m, 4H), 3.31-3.28 (m, 4H), 2.28 (s, 3H).

EXAMPLE 143

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0236] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (10 mg, 0.029 mmol) and 4-chlorobenzoyl chloride (6 mg, 0.035 mmol) similar to Example 139 and isolated as a white solid (7 mg, 50%). ¹H NMR (CDCl₃): 8.77 (d, *J* = 6.0 Hz, 2H), 8.29 (d, *J* = 6.0 Hz, 2H), 8.00 (s, 1H), 7.83 (d, *J* = 8.7 Hz, 2H), 7.81 (s, 1H), 7.49 (d, *J* = 8.4 Hz, 2H), 7.35 (s, 2H), 7.01 (s, 1H), 6.75 (s, 1H), 2.30 (2, 3H).

EXAMPLE 144

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0237] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (10 mg, 0.029 mmol) and p-anisoyl chloride (6 mg, 0.035 mmol) similar to Example 139 and isolated as a tan solid (10 mg, 71%). ¹H NMR (CDCl₃): 8.77 (dd, *J* = 1.5, 4.8 Hz, 2H), 8.29 (dd, *J* = 1.5, 4.8 Hz, 2H), 7.99 (s, 1H), 7.86 (d, *J* = 8.7 Hz, 2H), 7.79 (s, 1H), 7.36 (dd, *J* = 1.8, 8.4 Hz, 1H), 7.32 (d, *J* = 8.7 Hz, 2H), 7.02 (s, 1H), 6.99 (d, *J* = 8.7 Hz, 2H), 3.88 (s, 3H), 2.29 (s, 3H).

EXAMPLE 145

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0238] The title compound was prepared from a mixture of 4-(5-amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (80 mg, 0.232 mmol) and 4-chloromethylbenzoyl chloride (53 mg, 0.278 mmol) similar to Example 139 and isolated as a yellow solid (55 mg, 48%). ¹H NMR (CDCl₃): 8.75 (dd, *J* = 1.8, 4.8 Hz, 2H), 8.29 (dd, *J* = 1.8, 4.8 Hz, 2H), 8.02 (s, 1H), 7.96 (s, 1H), 7.88 (d, *J* = 8.4 Hz, 2H), 7.51 (d, *J* = 8.4 Hz, 2H), 7.37 (dd, *J* = 1.8, 7.5 Hz, 1H), 7.32 (d, *J* = 7.5 Hz, 1H), 7.12 (s, 1H), 6.75 (s, 1H), 4.63 (s, 2H), 2.28 (s, 3H).

EXAMPLE 146

4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide

[0239] To a stirring mixture of 4-chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide (48 mg, 0.096 mmol) and 1-methylpiperazine (16 μl, 0.0144 mmol) in anhydrous tetrahydrofuran (5 ml) was added N,N-diisopropylethylamine (200 μl). The mixture was refluxed for 20 h. The mixture was cooled to room temperature and then rotary evaporated to dryness. The residue was purified by column chromatography and isolated as a tan solid (22 mg, 41%). ¹H NMR (CDCl₃): 8.77 (dd, *J* = 1.8, 4.5 Hz, 2H), 8.29 (dd, *J* = 1.5, 4.5 Hz, 2H), 8.01 (s, 1H), 7.85 (s, 1H), 7.83 (d, *J* = 8.1 Hz, 2H), 7.47 (d, *J* = 7.8 Hz, 2H), 7.37 (dd, *J* = 1.5, 8.4 Hz, 1H), 7.33 (d, *J* = 8.4 Hz, 1H), 7.04 (s, 1H), 6.74 (s, 1H), 3.57 (s, 2H), 2.47 (s, 8H), 2.29 (s, 6H).

- 109 -

EXAMPLE 147

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0240] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-amino-2,5-dimethylphenol (40 mg, 0.290 mmol) similar to Example 117 and isolated as an orange solid (35 mg, 47%). ¹H NMR (CDCl₃): 8.79-8.77 (m, 1H), 8.58 (d, *J* = 7.8 Hz, 1H), 8.14 (bs, 1H), 7.91-7.86 (m, 1H), 7.46-7.42 (m, 1H), 7.09 (bs, 1H), 6.92 (s, 1H), 6.83 (s, 1H), 6.46 (s, 1H), 2.22 (s, 3H), 2.02 (s, 3H).

EXAMPLE 148

4-(2,5-Dimethyl-4-hydroxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0241] The title compound was prepared from a mixture of 4-chloro-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.191 mmol) and 4-amino-2,5-dimethylphenol (39 mg, 0.287 mmol) similar to Example 117 and isolated as an orange solid (35 mg, 47%). ¹H NMR (CDCl₃): 9.75 (d, *J* = 1.2 Hz, 1H), 8.79-8.78 (m, 1H), 8.73 (d, *J* = 2.4 Hz, 1H), 7.20 (s, 1H), 7.01 (s, 1H), 6.78 (s, 1H), 6.52 (s, 1H), 5.43 (s, 1H), 2.25 (s, 3H), 2.16 (s, 3H).

EXAMPLE 149

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0242] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-amino-2,5-dimethylphenol (40 mg, 0.290 mmol) similar to Example 117 and isolated as a white solid (2 mg). ¹H NMR (CDCl₃): 8.77 (d, *J* = 6.0 Hz, 2H), 8.27 (d, *J* = 6.3 Hz, 2H), 7.05 (s, 1H), 6.86 (s, 1H), 6.77 (s, 1H), 6.47 (s, 1H), 5.25 (bs, 1H), 2.26 (s, 3H), 2.19 (s, 3H).

- 110 -

EXAMPLE 150

4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0243] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-chloro-2,5-dimethoxyaniline (54 mg, 0.290 mmol) similar to Example 117 and isolated as a brown solid (58 mg, 73%). ¹H NMR (CDCl₃): 8.84-8.82 (m, 1H), 8.61-8.57 (m, 1H), 7.93-7.87 (m, 1H), 8.50 (bs, 1H), 7.66 (s, 1H), 7.49-7.45 (m, 1H), 7.02 (s, 1H), 7.01 (s, 1H), 4.07 (s, 3H), 3.92 (s, 3H).

EXAMPLE 151

4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0244] The title compound was prepared from a mixture of 4-chloro-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.191 mmol) and 4-chloro-2,5-dimethoxyaniline (54 mg, 0.287 mmol) similar to Example 117 and isolated as a yellow solid (67 mg, 85%). ¹H NMR (CDCl₃): 9.77 (d, *J* = 1.5 Hz, 1H), 8.79-8.78 (m, 1H), 8.76 (d, *J* = 2.4 Hz, 1H), 8.50 (bs, 1H), 7.70 (s, 1H), 7.05 (s, 1H), 7.04 (s, 1H), 4.06 (s, 3H), 3.93 (s, 3H).

EXAMPLE 152

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0245] The title compound was prepared from a mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4,5-dimethoxy-2-methylaniline (48 mg, 0.290 mmol) similar to Example 117 and isolated as a yellow solid (64 mg, 79%). ¹H NMR (CDCl₃): 8.82-8.80 (m, 1H), 8.55 (d, *J* = 7.8 Hz, 1H), 7.88-7.82 (m, 1H), 7.51 (s, 1H), 7.43-7.38 (m, 1H), 6.80 (s, 1H), 6.76 (s, 1H), 6.50 (s, 1H), 3.92 (s, 3H), 3.84 (s, 3H), 2.19 (s, 3H).

- 111 -

EXAMPLE 153

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0246] The title compound was prepared from a mixture of 4-chloro-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.191 mmol) and 4,5-dimethoxy-2-methylaniline (48 mg, 0.287 mmol) similar to Example 117 and isolated as a brown solid (47 mg, 56%). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.5 Hz, 1H), 8.76 (dd, *J* = 1.5, 2.4 Hz, 1H), 8.71 (d, *J* = 2.4 Hz, 1H), 7.51 (s, 1H), 6.82 (s, 1H), 6.78 (s, 1H), 6.55 (s, 1H), 3.93 (s, 3H), 3.86 (s, 3H), 2.21 (s, 3H).

EXAMPLE 154

4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0247] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4,5-dimethoxy-2-methylaniline (48 mg, 0.290 mmol) similar to Example 117 and isolated as a brown solid (23 mg, 30%). ¹H NMR (CDCl₃): 9.54 (dd, *J* = 1.5, 4.5 Hz, 2H), 9.05 (dd, *J* = 1.5, 4.5 Hz, 2H), 7.98 (s, 1H), 7.60 (s, 2H), 7.29 (s, 1H), 4.71 (s, 3H), 4.63 (s, 3H), 3.00 (s, 3H).

EXAMPLE 155

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-aminopyrimidine

[0248] The title compound was prepared from a mixture of 2-amino-4-chloro-6-methyl-pyrimidine (100 mg, 0.696 mmol) and 2-chloro-5-methoxyaniline hydrochloride (203 mg, 1.05 mmol) similar to Example 117 and isolated as a white solid (132 mg, 72%). ¹H NMR (CDCl₃): 7.77 (d, *J* = 2.7 Hz, 1H), 7.28 (d, *J* = 9.0 Hz, 6.74 (s, 1H), 6.57 (dd, *J* = 3.0, 8.7 Hz, 1H), 6.04 (s, 1H), 4.81 (s, 2H), 3.82 (s, 3H), 2.26 (s, 3H).

EXAMPLE 156

4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0249] A mixture of 4-chloro-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (25 mg, 0.096 mmol), 3-trifluoromethoxyaniline (19 μ l, 0.144 mmol), and 2N HCl (75 μ l) in water:ethanol (2:1, 5 ml) was refluxed for 24 h. The mixture was cooled to room temperature and the resulting precipitate was filtered, washed with water, water:ethanol (2:1) and dried to give a yellow solid (23 mg, 59%). ^1H NMR (CDCl_3): 8.88-8.85 (m, 1H), 8.59-8.55 (m, 1H), 7.93-7.87 (m, 1H), 7.59 (s, 1H), 7.48 (t, $J = 8.3$ Hz, 1H), 7.45 (dd, $J = 1.2, 7.5$ Hz, 1H), 7.36 (s, 1H), 7.30-7.29 (m, 1H), 7.16-7.10 (m, 1H), 7.05 (s, 1H).

EXAMPLE 157

4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0250] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4,5-dimethoxy-2-methylaniline (48 mg, 0.290 mmol) similar to Example 117 and isolated as a red solid (8 mg, 11%). ^1H NMR (CDCl_3): 9.69 (s, 1H), 8.74-8.68 (m, 2H), 7.56 (s, 1H), 7.43-7.39 (m, 1H), 6.84 (s, 1H), 6.83 (s, 1H), 6.46 (s, 1H), 3.94 (s, 3H), 3.86 (s, 3H), 2.23 (s, 3H).

EXAMPLE 158

4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride

[0251] The title compound was prepared from a mixture of 4-chloro-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-chloro-2,5-dimethoxyaniline (54 mg, 0.290 mmol) similar to Example 117 and isolated as a brown solid (55 mg, 68%). ^1H NMR (CDCl_3): 8.78 (dd, $J = 1.8$,

- 113 -

4.8 Hz, 2H), 8.31 (dd, $J = 1.8, 4.8$ Hz, 2H), 8.24 (s, 1H), 7.52 (s, 1H), 7.01 (s, 1H), 6.96 (s, 1H), 3.98 (s, 3H), 3.91 (s, 3H).

EXAMPLE 159

4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0252] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 2-hydroxy-5-methylaniline (36 mg, 0.290 mmol) similar to Example 117 and isolated as a brown solid (10 mg, 15%). ^1H NMR (Acetone- d_6): 9.58 (s, 1H), 8.87 (s, 1H), 8.74 (s, 1H), 8.72-8.68 (m, 2H), 7.89 (s, 1H), 7.58-7.53 (m, 1H), 7.24 (s, 1H), 6.92 (s, 2H), 2.33 (s, 3H).

EXAMPLE 160

4-(N-Methyl-3-methoxyanilino)-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine

[0253] A mixture of 4-chloro-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine (180 mg, 0.693 mmol) and N-methyl-3-methoxyaniline (112 mg, 0.816 mmol) in 3 mL of solvent (ethanol : water = 2:1) was refluxed for 20 h. The solvent was removed under reduced pressure and the residue was dissolved in 25 mL of ethyl acetate. The ethyl acetate solution was washed with 25 mL of 1M NaOH, and the aqueous layer was reextracted with ethyl acetate (25 mL). The combined organic extracts were washed with saturated NaCl and dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by flash column chromatography (40-45% ethyl acetate/hexanes) to give the title compound as an oil (250 mg, 0.693 mmol, 100%). ^1H NMR (CDCl_3): 8.86 (dd, $J = 0.6, 4.5$ Hz, 1H), 8.50 (d, $J = 7.5$ Hz, 1H), 7.85 (dt, $J = 1.8, 7.8$ Hz, 1H), 7.42 (m, 2H), 6.95 (dd, $J = 1.8, 8.4$ Hz, 1H), 6.86 (m, 1H), 6.82 (m, 1H), 3.85 (s, 3H), 3.68 (s, 3H).

- 114 -

EXAMPLE 161

4-(3,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine

[0254] The title compound was prepared from 4-chloro-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine (150 mg, 0.578 mmol) and 3,5-dimethoxyaniline (95 mg, 0.621 mmol) similar to Example 160 and was isolated as off-white solid (119 mg, 0.316 mmol, 55%). ¹H NMR (DMSO-*d*₆): 10.31 (s, br, 1H), 8.80 (m, 2H), 8.17 (m, 2H), 7.17 (s, 1H), 7.04 (s, 2H), 6.32 (m, 1H), 3.80 (s, 6H), 3.53 (s, 1H).

EXAMPLE 162

4-(5-Chloro-2-methoxyanilino)-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine

[0255] The title compound was prepared from 4-chloro-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine (150 mg, 0.578 mmol) and 5-chloro-2-methoxyaniline (98 mg, 0.621 mmol) similar to Example 160 and was isolated as off-white solid (84 mg, 0.221 mmol, 38%). ¹H NMR (CDCl₃/MeOH-*d*₄): 8.72 (d, *J* = 4.2 Hz, 2H), 8.51 (s, 1H), 8.34 (d, *J* = 4.5 Hz), 7.09 (m, 2H), 6.91 (d, *J* = 8.4 Hz, 1H), 3.93 (s, 3H)

EXAMPLE 163

4-(3,5-Dimethoxyanilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine

[0256] a. 4-Chloro-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine. A mixture of POCl₃ (freshly distilled, 0.50 mL, 5.35 mmol) in 2.5 mL of DMF was stirred at 0 °C for 45 min. To the mixture was added 2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-4-pyrimidinol (98 mg, 0.317 mmol) in 0.5 mL of DMF and the mixture was allowed to warm to room temperature. The resulting mixture was heated overnight at 100 °C. The reaction mixture was cooled to room temperature and it was extracted with ethyl acetate (10 mL X 3). The combined extracts were washed with water (10 mL X 3) and

- 115 -

saturated NaCl, dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash column chromatography (5% ethyl acetate/hexanes) to obtain the title compound as a white solid (74 mg, 0.226 mmol, 72%, mp 109-110 °C). ¹H NMR (CDCl₃): 8.30 (s, 2H), 7.51 (s, 1H).

[0257] b. 4-(3,5-Dimethoxyanilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine. The title compound was prepared from 4-chloro-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine (57 mg, 0.174 mmol) and 3,5-dimethoxyaniline (38 mg, 0.248 mmol) similar to Example 160 and was isolated as a yellow crystalline solid (58 mg, 0.130 mmol, 75%). ¹H NMR (CDCl₃): 8.25 (s, 2H), 7.25 (s, br, 1H), 7.02 (s, 1H), 6.62 (m, 2H), 6.38 (t, *J* = 1.8 Hz, 1H), 3.84 (s, 6H).

EXAMPLE 164

4-(2-Chloro-5-methoxyanilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine

[0258] The title compound was prepared from 4-chloro-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine (52 mg, 0.159 mmol) and 2-chloro-5-methoxyaniline (40 mg, 0.255 mmol) similar to Example 160 and was isolated as a yellow crystalline solid (13 mg, 0.029 mmol, 18%). ¹H NMR (CDCl₃): 8.27 (s, 2H), 7.76 (s, br, 1H), 7.40 (s, 1H), 7.38 (s, 1H), 7.02 (s, 1H), 6.77 (m, 1H), 3.90 (s, 3H).

EXAMPLE 165

4-(2-Methoxy-5-methyl-4-nitroanilino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0259] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (51 mg, 0.194 mmol) and 2-methoxy-5-methyl-4-nitroaniline (43 mg, 0.236 mmol) similar to Example 160 and was isolated as off-white crystalline solid (mp 128-133 °C, 23 mg, 0.057 mmol, 29%). ¹H NMR (CDCl₃/MeOH-*d*₄): 9.42 (s, 1H), 8.75 (m, 1H), 8.62 (m, 1H), 7.77 (s,

- 116 -

1H), 7.68 (s, 1H), 7.42 (dd, $J = 4.8, 8.1$ Hz, 1H), 6.82 (s, 1H), 4.01 (s, 3H), 2.01 (s, 3H).

EXAMPLE 166

4-(1H-5-Indolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0260] A mixture of 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (42 mg, 0.162 mmol), 1H-5-indolylamine (31 mg, 235 mmol) and potassium carbonate (25 mg, 0.180 mmol) was heated overnight in 1 mL of DMF at 100 °C in a sealed tube. The reaction mixture was cooled to room temperature, diluted with 20 mL of ethyl acetate, washed with water (20 mL X 3) and saturated NaCl. The organic layer was dried over Na₂SO₄, filtered, concentrated and the residue was purified by flash column chromatography (30% ethyl acetate/hexanes) to obtain the title compound as a white solid (mp 142-143 °C, 45 mg, 0.127 mmol, 78%). ¹H NMR (CDCl₃): 9.56 (d, $J = 1.5$ Hz, 1H), 8.74 (dt, $J = 1.8, 8.1$ Hz, 1H), 8.63 (dd, $J = 3, 5.1$ Hz, 1H), 7.66 (m, 1H), 7.46 (m, 2H), 7.30 (d, $J = 3$ Hz, 1H), 7.19 (m, 1H), 6.82 (s, 1H), 6.55 (d, $J = 3$ Hz, 1H).

EXAMPLE 167

4-(5-Methoxy-2-methyl-4-nitroanilino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0261] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (48 mg, 0.185 mmol) and 5-methoxy-2-methyl-4-nitroaniline (35 mg, 0.192 mmol) similar to Example 166 and was isolated as a white solid (mp 128-132 °C, 11 mg, 0.027 mmol, 15%). ¹H NMR (CDCl₃): 9.62 (s, 1H), 8.68 (m, 2H), 7.79 (s, 1H), 7.73 (s, 1H), 7.48 (m, 1H), 7.01 (s, 1H), 4.02 (s, 3H), 2.63 (s, 1H).

- 117 -

EXAMPLE 168

4-(3-Trifluoromethyl-1-pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0262] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (74 mg, 0.285 mmol) and 3-trifluoromethyl-pyrazol (48 mg, 0.353 mmol) similar to Example 166 and was isolated as a white solid (mp 139-140 °C, 53 mg, 0.148 mmol, 52%). ¹H NMR (CDCl₃): 9.73 (s, 1H), 8.82 (m, 3H), 8.26 (s, 1H), 7.50 (m, 1H), 6.87 (d, *J* = 2.7 Hz, 1H).

EXAMPLE 169

4-(4,5-Dihydro-2-thiazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0263] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (75 mg, 0.289 mmol) and 2-amino-2-thiazoline (37 mg, 0.362 mmol) similar to Example 166 and was isolated as a white solid (20 mg, 0.61 mmol, 21%). ¹H NMR (CDCl₃/MeOH-*d*₄) 9.45 (s, 1H), 8.73 (dt, 1.8, 8.1 Hz, 1H), 8.58 (dd, *J* = 1.8, 5.1 Hz, 1H), 7.47 (m, 1H), 6.70 (s, 1H), 3.93 (t, *J* = 6.6 Hz, 2H), 3.04 (t, *J* = 7.2 Hz, 2H).

EXAMPLE 170

4-(1-Pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0264] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (74 mg, 0.285 mmol) and pyrazole (28 mg, 0.425 mmol) similar to Example 166 and was isolated as a white solid (mp 151-152 °C, 67 mg, 0.230 mmol, 81%). ¹H NMR (CDCl₃) 9.71 (s, 1H), 8.77 (m, 3H), 8.19 (s, 1H), 7.88 (m, 1H), 7.47 (m, 1H), 6.61 (dd, *J* = 1.5, 2.7 Hz, 1H).

- 118 -

EXAMPLE 171

4-(4,5,6,7-Tetrahydro-1H-indazolyl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0265] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (76 mg, 0.293 mmol) and 4,5,6,7-tetrahydroindazole (44 mg, 0.360 mmol) similar to Example 166 and was isolated as a white solid (25 mg, 0.072 mmol, 25% %). ¹H NMR (CDCl₃) 9.68 (d, *J* = 1.2 Hz, 1H), 8.75 (m, 2H), 8.40 (s, 1H), 8.07 (s, 1H), 7.45 (ddd, *J* = 1.2, 5.1, 9.0 Hz, 1H), 2.73 (m, 4H), 1.88 (m, 4H).

EXAMPLE 172

4-(1H-3-Pyrazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine

[0266] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine (75 mg, 0.289 mmol) and 3-amino pyrazole (31 mg, 0.373 mmol) similar to Example 166 and was isolated as a white solid (mp 230 °C decompose, 31 mg, 0.101 mmol, 35 %). ¹H NMR (DMSO-*d*₆) 9.62 (d, *J* = 1.2 Hz, 1H), 8.77 (m, 3H), 7.71 (s, 1H), 7.62 (dd, *J* = 4.5, 7.5 Hz, 1H), 6.09 (d, *J* = 3 Hz, 1H), 5.89 (s, 1H).

EXAMPLE 173

6-Methyl-2-(2-pyridinyl)-4-(3-trifluoromethylbenzylamino)-pyrimidine

[0267] A mixture of 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol), 3-trifluoromethylbenzylamine (70.1 mg, 0.4 mmol), water (5 ml) and 2N HCl (300 μl) was refluxed for two days. The resulting solution was basified to pH ~ 11 with ammonia and extracted with ethyl acetate (10 ml X 3). The combined ethyl acetate extracts were dried with sodium sulfate, and evaporated under vacuum. The residue was purified by column chromatography (hexane/ethyl acetate: 1/1) to give the title compound as a

- 119 -

solid (34 mg, 25 %). ^1H NMR (CD_3OD): 8.59 (d, $J = 4.2$ Hz, 1H), 8.27 (m, 2H), 7.86 (m, 2 H), 7.43 (m, 2H), 6.18 (s, 2H), 4.76 (s, 2H), 2.26 (s, 3H).

EXAMPLE 174

6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine

[0268] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol) and 3-phenoxyaniline (74.0 mg, 0.4 mmol) similar to Example 173 and was isolated as a white solid. (96 mg, 68 %). ^1H NMR (CDCl_3): 8.62 (d, $J = 4.2$ Hz, 1H), 8.25 (d, $J = 8.1$ Hz, 1H), 7.65 (t, $J = 1.5$ Hz, 1H), 7.21 (m, 5H), 6.94 (m, 5H), 6.64 (d, $J = 8.1$ Hz, 1H), 6.49 (s, 1H), 2.33 (s, 3H),

EXAMPLE 175

4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0269] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol) and 3-chloroaniline (50.8 mg, 0.4 mmol) similar to Example 173 and was isolated as a white solid (99 mg, 84 %). ^1H NMR (CDCl_3): 8.59 (d, $J = 4.2$ Hz, 1H), 8.29 (d, $J = 8.1$ Hz, 1H), 7.80 (br, 1H), 7.67 (t, $J = 1.5$ Hz, 1H), 7.38 (s, 1H), 7.19 (m, 1H), 7.10 (m, 2H), 6.91 (m, 1H), 6.41(s, 1H), 2.28 (s, 3H).

EXAMPLE 176

4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0270] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol) and 3,4-dimethoxyaniline (61.2 mg, 0.4 mmol) similar to Example 173 and was isolated as a white solid (108 mg, 84 %). ^1H NMR (CDCl_3): 8.63 (d, $J = 4.2$ Hz, 1H), 8.29 (d, $J = 8.1$ Hz, 1H),

- 120 -

7.65 (t, $J = 1.5$ Hz, 1H), 7.18 (m, 2 H), 6.7 (s, 1H), 6.71 (s, 2H), 6.28 (s, 1H), 3.73 (s, 3H), 3.71 (s, 3H), 2.29 (s, 3H).

EXAMPLE 177

4-(4-Fluoro-3-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0271] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol) and 4-fluoro-3-methoxyaniline (56.4 mg, 0.4 mmol) similar to Example 173 and was isolated as a white solid (96.7 mg, 78 %). ^1H NMR (CDCl_3): 8.63 (d, $J = 4.2$ Hz, 1H), 8.29 (d, $J = 7.5$ Hz, 1H), 7.65 (t, $J = 7.5$ Hz, 1H), 7.42 (s, 1H), 7.20 (m, 1H), 7.0 (d, $J = 7.5$ Hz, 1H), 6.90 (m, 1H), 6.69 (m, 1H), 3.71 (s, 3H), 2.30 (s, 3H),

EXAMPLE 178

4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine

[0272] The title compound was prepared from 4-chloro-6-methyl-2-(2-pyridinyl)pyrimidine (82.3 mg, 0.4 mmol) and 3-isopropoxyaniline (60.4 mg, 0.4 mmol) similar to Example 173 and was isolated as a white solid (88 mg, 69 %). ^1H NMR (CDCl_3): 8.64 (d, $J = 4.2$ Hz, 1H), 8.35 (d, $J = 8.1$ Hz, 1H), 7.7 (t, $J = 8.1$ Hz, 1H), 7.24 (m, 1H), 7.15 (m, 1H), 6.96 (s, 1H), 6.76 (m, 2H), 6.60 (d, $J = 8.1$ Hz, 1H), 6.18 (m, 1H), 4.45 (m, 1H), 2.37 (s, 3H), 1.25 (d, $J = 6.0$ Hz, 6H).

EXAMPLE 179

4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-pyridinyl)pyrimidine

[0273] a. 4-Hydroxy-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine. To a solution of pyridine-2-carboxamide hydrochloride (5.0 g, 33.2 mmol) in 20 ml of water was added concentrated aqueous KOH and the resulting mixture was extracted with ethyl acetate (50 mL X 3). The extract was evaporated

- 121 -

under vacuum to afford 4.0 g of free base. To a solution of the free base and 4,4,4-trifluoro-but-2-ynoic acid ethyl ester (6.0 g, 36 mmol) in ethanol (50 mL) at 0 °C ~ 60 °C was added dropwise potassium hydroxide solution (1.68 g in 40 mL water) in twenty minutes. The solution was then refluxed 2.5 h. The solvent was evaporated under vacuum and the residue was extracted with ethyl acetate (30 ml X 3). The combined ethyl acetate extracts were dried with sodium sulfate, evaporated to give a solid. It was recrystallized with Hexane/methylene chloride/methanol: 6/2/1) to give the product as a white solid (2.50 g, 31.2 %). ¹H NMR (CD₃OD): 8.63(d, *J* = 4.8 Hz, 1H), 8.31(d, *J* = 7.8 Hz, 1H), 7.90 (t, *J* = 7.8 Hz, 1H), 7.49 (m, 1H), 6.73 (s, 1H), 4.76 (s, 1H).

[0274] b. 4-Chloro-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine. A mixture of 4-hydroxy-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine (2.50 g, 10.37 mmol) and phosphorus oxychloride (20 mL) was refluxed for 3 h. The resulting solution was evaporated under vacuum and the residue was basified to pH ~ 11 with ammonia and extracted with ethyl acetate (15 ml X 3). The combined ethyl acetate extracts were dried with sodium sulfate, evaporated to give a solid. The solid was purified by column chromatography (hexane/ethyl acetate: 2/1 ~ 1/3) to afford a yellow solid (2.30 g, 86 %). ¹H NMR (CDCl₃): 8.76 (d, *J* = 2.7 Hz, 1H), 8.42(d, *J* = 7.8 Hz, 1H), 7.76 (t, *J* = 1.8 Hz, 1H), 7.55 (d, *J* = 2.4 Hz, 1H), 7.34 (t, *J* = 5.1Hz, 1H).

[0275] c. 4-(3,4-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine. The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine (51.8 mg, 0.2 mmol) and 3,4-dimethoxyaniline (51.8 mg, 0.2 mmol) similar to Example 173 and was isolated as a white solid (61 mg, 81%). ¹H NMR (CDCl₃): 8.71 (d, *J* = 0.9 Hz, 1H), 8.43 (d, *J* = 7.8 Hz, 1H), 7.45 (t, *J* = 7.8 Hz, 1H), 7.50 (s, 1H), 7.35 (m, 1H), 6.77 (m, 4H), 3.80 (s, 3H), 3.77 (s, 3H).

EXAMPLE 180

4-(5-Chloro-2-methoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0276] A mixture of 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (100 mg, 0.385 mmol), 2-methoxy-5-chloroaniline (90 mg, 0.577 mmol) and 2N HCl (150 μ l) in 1:1 ethanol/water (5 ml) was refluxed for 48 h. The resulting yellow solid was collected by filtration, washed with water, ethanol/water (1:4) and then dried under vacuum to give the title compound (100 mg, 67%). ^1H NMR (CDCl_3): 9.66 (d, $J = 1.8$ Hz, 1H), 8.70-8.76 (m, 2H), 7.68 (dd, $J = 7.5, 1.0$ Hz, 1H), 7.44 (dd, $J = 7.5, 1.0$ Hz, 1H), 7.09 (dd, $J = 8.7, 2.7$ Hz, 1H), 6.92 (s, 1H), 6.87 (d, $J = 8.7$ Hz, 1H), 3.91 (s, 3H).

EXAMPLE 181

4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0277] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.153 mmol) and 2-methoxy-5-nitroaniline (39 mg, 0.230 mmol), similar to Example 180 and was isolated as a yellow solid (30 mg, 51%). ^1H NMR (CDCl_3): 9.760 (d, $J = 1.5$ Hz, 1H), 8.86 (dd, $J = 1.8, 2.7$ Hz, 1H), 8.74 (d, $J = 2.4$ Hz, 1H), 8.10 (dd, $J = 2.7, 9.0$ Hz, 1H), 7.34 (s, 1H), 7.07 (d, $J = 9.0$ Hz, 1H), 4.08 (s, 3H).

EXAMPLE 182

4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0278] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.153 mmol) and 5-methoxy-2-nitroaniline (39 mg, 0.230 mmol), similar to Example 180 and was isolated as an off white solid (35 mg, 58%). ^1H NMR (CDCl_3): 9.69 (d, $J = 1.5$ Hz, 1H), 8.83 (dd, $J = 1.5, 2.4$ Hz, 1H), 8.72 (d, $J = 2.4$ Hz, 1H), 8.08 (d, $J = 9.6$ Hz, 1H), 7.11 (s, 1H), 6.27 (dd, $J = 2.7, 9.6$ Hz, 1H), 6.15 (d, $J = 2.7$ Hz, 1H), 3.83 (s, 3H).

EXAMPLE 183

4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0279] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.153 mmol) and 3,5-dimethoxyaniline (35 mg, 0.23 mmol), similar to Example 180 and was isolated as a pale solid (30 mg, 52%). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.5 Hz, 1H), 7.78 (m, 1H), 8.72 (d, *J* = 2.4 Hz, 1H), 7.48 (s, 1H), 7.10 (s, 1H), 6.53 (d, *J* = 2.1 Hz, 2H) 6.39 (t, *J* = 2.1 Hz, 1H), 3.83 (s, 6H).

EXAMPLE 184

4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0280] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 5-carboxy-2-methoxyaniline (38 mg, 0.231 mmol), similar to Example 180 and was isolated as an off white solid (20 mg, 33%). ¹H NMR (CDCl₃): 9.99 (brs, 1H), 9.660 (d, *J* = 1.5 Hz, 1H), 8.86 (m, 3H), 7.80 (d, 2H), 7.2 (m, 2 H), 3.94 (s, 3H).

EXAMPLE 185

4-(5-Hydroxy-2-nitroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0281] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (60 mg, 0.23 mmol) and 5-hydroxy-2-nitroaniline (52 mg, 0.34 mmol) similar to Example 180. The reaction mixture was neutralized with 2N NaOH to pH 7.0 and the solvent was removed under vacuo. The product was purified by column chromatography (20% ethyl acetate in hexane) to give 55 mg (63%) of the title compound as a light yellow solid. ¹H NMR (CDCl₃): 9.40 (d, *J* = 1.2 Hz, 1H), 8.71 (dd, *J* = 1.2, 4.8 Hz, 1H), 8.57(ddd, *J* = 2.1, 3.9, 7.8 Hz, 1H), 8.04 (d, *J* = 2.7 Hz, 1H), 7.42 (dd, *J*

- 124 -

= 3.9, 8.7 Hz, 1H), 7.32 (dd, J = 6.0, 9.0 Hz, 1H), 7.23 (s, 1H), 6.92 (d, J = 9.0 Hz, 1H), 6.19 (s, 2H).

EXAMPLE 186

4-(2-Ethylsulfonyl-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0282] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2-ethylsulfonyl-5-hydroxyaniline (46 mg, 0.231 mmol) similar to Example 180. The reaction mixture was evaporated under vacuum to give a residue. The residue was precipitated using acetone to give the product as a yellow solid (30 mg, 46%). ^1H NMR (DMSO- d_6): 9.45 (s, 1H), 8.71 (m, 3H), 7.84 (brs, 1H), 7.0 (brd, 1H), 6.08 (brd, 1H), 3.0 (q, 2H), 1.0 (t, 3H).

EXAMPLE 187

4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0283] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.152 mmol) and 2-methoxy-5-methylaniline (28 mg, 0.231 mmol) similar to example 180 and was isolated as a solid (43 mg, 81%). ^1H NMR (CDCl_3): 9.73 (m, 1H), 8.44-8.41 (m, 1H), 7.81-7.75 (m, 1H), 7.46 (s, 1H), 7.34-7.28 (m, 2H), 7.23 (t, J = 8.1 Hz, 1H), 7.09-7.06 (m, 1H), 7.00-6.97 (m, 1H), 6.55 (d, J = 0.6 Hz, 1H), 2.44 (s, 3H), 2.43 (d, J = 0.6, 3H).

EXAMPLE 188

4-(5-Hydroxy-2-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

[0284] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 5-hydroxy-2-methylaniline (28 mg, 0.231 mmol) similar to example 185 and was isolated

- 125 -

as a pale solid (40 mg, 75%). ^1H NMR (CDCl_3): 9.73 (m, 1H), 8.44-8.41 (m, 1H), 7.81-7.75 (m, 1H), 7.46 (s, 1H), 7.34-7.28 (m, 2H), 7.23 (t, $J = 8.1$ Hz, 1H), 7.09-7.06 (m, 1H), 7.00-6.97 (m, 1H), 6.55 (d, $J = 0.6$ Hz, 1H), 2.44 (s, 3H), 2.43 (d, $J = 0.6$, 3H).

EXAMPLE 189

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluormethylpyrimidine

[0285] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2-chloro-5-hydroxyaniline (30 mg, 0.231 mmol) similar to Example 185 and was isolated as an off white solid (15 mg, 28%). ^1H NMR ($\text{DMSO}-d_6$): 9.89 (s, 1H), 9.85 (brs, NH), 8.74(d, $J = 5.7$, 1H), 8.22 (d, $J = 7.8$ Hz, 1H), 7.97 (ddd, $J = 1.8$, 5.7, 7.8 Hz, 1H), 7.54 (ddd, $J = 4.5$, 6.0, 7.8 Hz, 1H), 7.35 (m, 2H), 7.19 (s, 1H), 6.70 (dd, $J = 2.7$, 5.7 Hz, 1H).

EXAMPLE 190

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

[0286] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (60 mg, 0.23 mmol) and 2-methoxy-5-methylaniline (46 mg, 0.34 mmol) similar to Example 180. After completion of the reaction the solvent was removed under reduced pressure. The product was purified (5 mg) by preparative TLC plate using 1:1 hexane/ethyl acetate as mobile phase. ^1H NMR ($\text{DMSO}-d_6$): 9.68 (s, 1H), 8.75(d, $J = 3.6$, 1H), 8.26 (d, $J = 7.2$ Hz, 1H), 8.00 (t, $J = 7.2$ Hz, 1H), 7.55 (dd, $J = 4.0$, 7.2 Hz, 1H), 7.04 (m, 2H), 3.82 (s, 3H), 2.02 (s, 3H).

- 126 -

EXAMPLE 191

4-(2-Ethylsulfonyl-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

[0287] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2-ethylsulfonyl-5-hydroxyaniline (46 mg, 0.231 mmol) similar to Example 186 and was isolated as a yellow solid (35 mg, 54%). ¹H NMR (CDCl₃): 8.88 (s, 1H), 8.60 (brt, 2H), 8.01 (t, 1H), 7.52 (m, 2H), 7.01 (m, 2H), 3.02 (q, 2H), 1.22 (t, 3H).

EXAMPLE 192

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

[0288] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2-hydroxy-5-isopropylaniline (35 mg, 0.231 mmol) similar to Example 185 and was isolated as a solid (32 mg, 55%). ¹H NMR (DMSO-d₆): 8.74 (s, 1H), 8.73 (d, *J* = 0.9 Hz, 1H), 8.35 (d, *J* = 7.8 Hz, 2H), 7.95 (t, *J* = 8.1 Hz, 2H), 7.54 (m, 2H), 6.60 (m, 2H), 2.78 (sep, *J* = 6.9 Hz, 1H), 1.20 (d, *J* = 6.9 Hz, 6H).

EXAMPLE 193

4-(5-Hydroxy-2-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0289] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 5-hydroxy-2-methylaniline (34 mg, 0.231 mmol) similar to Example 190 and was isolated as a solid (3 mg). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.2 Hz, 1H), 8.76-8.71 (m, 2H), 7.30 (brs, 1H), 7.18 (d, *J* = 8.4 Hz, 1H), 6.87 (brd, 1H), 6.78 (dd, *J* = 8.4, 2.4 Hz, 1H), 6.73 (s, 1H), 2.21 (s, 3H).

- 127 -

EXAMPLE 194

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0290] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2-chloro-5-hydroxyaniline (33 mg, 0.231 mmol) similar to Example 190 and was isolated as a solid (6 mg). ¹H NMR (CDCl₃): 9.72 (d, *J* = 1.5 Hz, 1H), 8.78 (dd, *J* = 5.7, 1.5 Hz, 1H), 8.73 (d, *J* = 1.2 Hz, 1H), 7.33 (d, *J* = 8.7 Hz, 1H), 7.05 (s, 1H), 7.04 (d, *J* = 8.7 Hz, 1H), 6.73 (dd, *J* = 8.7, 2.7 Hz, 1H), 6.29 (d, *J* = 2.7 Hz, 1H), 6.19 (dd, *J* = 8.4, 2.7 Hz, 1H).

EXAMPLE 195

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

[0291] The title compound was prepared from 4-chloro-2-(2-pyridinyl)-6-trifluoromethylpyrimidine (30 mg, 0.115 mmol) and 3,5-dimethoxyaniline (26 mg, 0.172 mmol) similar to Example 180 and was isolated as a light yellow solid (16 mg, 37% yield). ¹H NMR (CDCl₃): 8.85 (dd, *J* = 4.8, 1.5 Hz, 1H), 8.56 (d, *J* = 6.9 Hz, 1H), 7.89 (ddd, *J* = 9.3, 7.5, 1.5 Hz, 1H), 7.65 (brs, 1H), 7.45 (dd, *J* = 9.3, 4.5 Hz, 1H), 7.09 (s, 1H), 6.51 (d, *J* = 1.8 Hz, 1H), 6.37 (t, *J* = 1.8 Hz, 1H), 3.93 (s, 6H).

EXAMPLE 196

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0292] The title compound was prepared similar to 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (30 mg, 0.115 mmol) and 2,5-dimethylaniline (21 mg, 0.172 mmol) similar to Example 180 and was isolated as an off white solid (35 mg, 80% yield). ¹H NMR (CDCl₃): 9.74 (d, *J* = 1.5 Hz, 1H), 8.76 (dd, *J* = 1.5, 2.4 Hz, 1H), 8.71 (d, *J* = 2.4 Hz, 1H), 7.48 (brs, 1H), 7.22 (d, *J* = 8.1 Hz, 1H), 7.12 (m, 2H), 6.65 (s, 1H), 2.36 (s, 3H), 2.23 (s, 3H).

EXAMPLE 197

4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0293] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (30 mg, 0.115 mmol) and 5-chloro-2-methoxyaniline (27 mg, 0.173 mmol) similar to Example 185 and was isolated as a solid (10 mg, 23% yield). ¹H NMR (CDCl₃): 9.72 (d, *J* = 1.2 Hz, 1H), 8.82 (t, *J* = 1.8 Hz, 1H), 8.72 (d, *J* = 2.4 Hz, 1H), 8.23 (brs, 1H), 7.71 (brs, 1H), 7.15 (dd, *J* = 8.4, 2.4 Hz, 1H), 7.04 (s, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), 3.90 (s, 3H).

EXAMPLE 198

4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0294] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (30 mg, 0.115 mmol) and 5-chloro-2-hydroxyaniline (25 mg, 0.173 mmol) similar to Example 185 and was isolated as a solid (7 mg, 16% yield). ¹H NMR (CDCl₃): 9.68 (d, *J* = 0.9 Hz, 1H), 8.84 (t, *J* = 2.4 Hz, 1H), 8.60 (d, *J* = 2.4 Hz, 1H), 7.89 (brs, 1H), 7.41 (s, 1H), 7.07 (m, 2H), 6.91 (d, *J* = 8.7 Hz, 1H).

EXAMPLE 199

4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine

[0295] The title compound was prepared from 4-chloro-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine (30 mg, 0.115 mmol) and 5-hydroxy-2-isopropylaniline (26 mg, 0.173 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.75 (d, *J* = 1.2 Hz, 1H), 8.77 (t, *J* = 1.8 Hz, 1H), 8.72 (d, *J* = 2.7 Hz, 1H), 7.29 (brs, 1H), 7.27 (s, 1H), 6.91 (dd, *J* = 8.4, 2.4 Hz, 1H), 6.79 (d, *J* = 2.7 Hz, 1H), 6.70 (s, 1H), 4.13 (sep, *J* = 6.9 Hz, 1H), 1.20 (d, *J* = 6.9 Hz, 6H).

- 129 -

EXAMPLE 200

4-(2,5-Dimethoxyphenylethylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0296] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (40 mg, 0.154 mmol) and 2,5-dimethoxyphenylethylamine (28 mg, 0.232 mmol) similar to Example 186 and was isolated as an off white solid (35 mg, 54% yield). ¹H NMR (CDCl₃): 9.59 (brs, 1H), 8.69 (d, *J* = 4.8 Hz, 1H), 7.37 (dd, *J* = 4.8, 8.4 Hz, 1H), 6.76 (m, 3H), 3.95 (s, 3H), 3.90 (s, 3H), 3.00 (t, 2H), 1.67 (s, 2H).

EXAMPLE 201

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0297] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 2,5-dimethyl-4-hydroxyaniline (40 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.51 (brs, 1H), 8.72 (d, *J* = 8.4 Hz, 1H), 8.63 (d, *J* = 3.6 Hz, 1H), 7.45 (dd, *J* = 7.5, 3.5 Hz, 1H), 7.33 (s, 1H), 7.00 (s, 1H), 6.74 (s, 1H), 6.45 (brs, 1H), 2.25 (s, 3H), 2.16 (s, 3H).

EXAMPLE 202

4-(3,4,5-Trichloroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0298] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 3,4,5-trichloroaniline (56 mg, 0.288 mmol) similar to Example 180 and was isolated as a white solid (30 mg, 37%). ¹H NMR (CDCl₃): 9.58 (d, *J* = 1.5 Hz, 1H), 8.99 (ddd, *J* = 8.4, 4.9, 1.5 Hz, 1H), 8.79 (dd, *J* = 5.4, 1.5 Hz, 1H), 7.99 (s, 1H), 7.76 (dd, *J* = 7.8, 5.1 Hz, 1H), 7.40 (s, 1H).

- 130 -

EXAMPLE 203

4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0299] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 2-cyano-4,5-dimethoxyaniline (48 mg, 0.288 mmol) similar to Example 185 and isolated as a solid (8 mg, 10%). ¹H NMR (CDCl₃): 9.62 (d, *J* = 1.4 Hz, 1H), 8.69 (m, 2H), 7.43 (dd, *J* = 5.4, 1.5 Hz, 1H), 7.07 (s, 1H), 6.93 (s, 1H), 6.70 (s, 1H).

EXAMPLE 204

4-(3-Methoxy-dibenzofuran-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0300] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 4-amino-3-methoxydibenzofuran (61 mg, 0.288 mmol) similar to Example 185 and was isolated as a solid (6 mg, 7%). ¹H NMR (CDCl₃): 9.73 (brs, 1H), 8.88 (d, *J* = 7.8 Hz, 1H), 8.77 (brs, 1H), 7.89 (m, 1H), 7.19-7.59 (m, 6H), 7.02 (s, 1H), 4.05 (s, 3H).

EXAMPLE 205

4-(1,5,6-Trimethyl-benzimidazol-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0301] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 1,5,6-trimethyl-4-aminobenzimidazole (50 mg, 0.288 mmol) similar to Example 185 and was isolated as a solid (5 mg, 6%). ¹H NMR (CDCl₃): 9.36 (ddd, *J* = 7.8, 3.0, 1.2 Hz, 1H), 9.68 (s, 1H), 9.01 (ddd, *J* = 5.7, 2.7, 1.2 Hz, 1H), 8.87 (s, 1H), 8.46 (dd, *J* = 8.4, 6.0 Hz, 1H), 8.04 (s, 1H), 7.71 (s, 1H), 7.48 (s, 1H), 6.93 (s, 1H), 2.62 (s, 3H), 2.24 (s, 3H), 2.21 (s, 3H).

- 131 -

EXAMPLE 206

4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0302] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 3-amino-2,6-dimethoxypyridine (44 mg, 0.288 mmol) similar to Example 180 and was isolated as a yellow solid (15 mg, 20%). ¹H NMR (CDCl₃): 9.62 (brs, 1H), 8.72-8.67 (m, 2H), 7.42 (dd, *J* = 8.1, 5.4 Hz, 1H), 7.09 (s, 1H), 6.75 (s, 1H), 6.43 (d, *J* = 8.7 Hz, 1H), 4.08 (s, 3H), 4.02 (s, 3H).

EXAMPLE 207

4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0303] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 5-amino-2-methoxypyridine (35 mg, 0.288 mmol) similar to Example 180 and was isolated as a solid (30 mg, 45%). ¹H NMR (CDCl₃): 9.65 (brs, 1H), 8.72-8.69 (m, 2H), 8.26 (d, *J* = 2.7 Hz, 1H), 7.71 (d, *J* = 6.9 Hz, 1H), 7.44-7.40 (m, 2H), 6.86 (d, *J* = 9.0 Hz, 1H), 6.74 (s, 1H), 3.98 (s, 3H).

EXAMPLE 208

4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0304] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 2-amino-4,6-dimethoxypyrimidine (27 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.65 (brs, 1H), 8.71-8.70 (m, 2H), 7.54 (brs, 1H), 7.44 (dd, *J* = 7.8, 4.8 Hz, 1H), 7.25 (s, 1H), 7.04 (s, 1H), 6.07 (t, *J* = 5.4 Hz, 1H).

- 132 -

EXAMPLE 209

4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0305] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 4-amino-5-chloro-2-methylpyrimidine (42 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.58 (d, *J* = 1.5 Hz, 1H), 8.72-8.65 (m, 2H), 7.42 (dd, *J* = 6, 5.1 Hz, 1H), 6.70 (s, 1H), 5.32 (s, 2H), 1.70 (s, 3H).

EXAMPLE 210

4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0306] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 4-amino-2-hydroxy-5-methylpyrimidine (36 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.47 (brs, 1H), 8.59-8.54 (m, 2H), 7.28 (dd, *J* = 8.1, 5.1 Hz, 1H), 6.59 (s, 1H), 5.18 (s, 2H), 1.70 (s, 3H).

EXAMPLE 211

4-(1,3,4-Triazol-1-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

[0307] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 1-amino-1,3,4-triazole (24 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.61 (brs, 1H), 8.71-8.66 (m, 4H), 7.41 (m, 2H), 6.69 (s, 1H), 5.18 (s, 2H).

- 133 -

EXAMPLE 212

4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine

- [0308] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 4-aminoindole (37 mg, 0.288 mmol) similar to Example 180 and was isolated as a solid (55 mg, 80%). ¹H NMR (CDCl₃): 8.76 (brs, 1H), 8.58 (d, *J* = 8.1 Hz, 1H), 7.93 (t, *J* = 7.5 Hz, 1H), 7.51-7.39 (m, 3H), 7.28-7.16 (m, 3H), 6.94 (s, 1H), 6.45 (s, 1H).

EXAMPLE 213

4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine

- [0309] The title compound was prepared from 4-chloro-2-(3-pyridinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 5-amino-2-methylindole (41 mg, 0.288 mmol) similar to Example 190 and was isolated as a solid. ¹H NMR (CDCl₃): 9.66 (s, 1H), 8.75-8.72 (m, 2H), 8.08 (s, 1H), 7.46-7.34 (m, 4H), 7.10 (d, *J* = 8.4 Hz, 1H), 6.78 (s, 1H), 6.45 (s, 1H).

EXAMPLE 214

4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride

- [0310] a. 2-Amidinopyrimidine: A solution of 2-cyanopyrimidine (4 g, 38.06 mmol) in 7N ammonia in methanol (300 ml) was stirred at room temperature for 48 h. The mixture was rotary evaporated to dryness. The residue was washed with hexane, filtered, and dried under vacuo to give a pale white solid (3.5 g, 76%). ¹H NMR (DMSO-d₆): 8.96 (d, *J* = 4.8 Hz, 2H), 7.63 (t, *J* = 4.8 Hz, 1H), 7.10 (bs, 3H).
- [0311] b. 2-(2-Pyrimidinyl)-6-trifluoromethylpyrimidin-4-ol: To a stirring solution of ethanolic sodium ethoxide (100 mg of sodium ethoxide in 20 ml of ethanol) was added 2-amidinopyrimidine (500 mg, 4.09 mmol) and

- 134 -

ethyl 4,4,4-trifluoroacetoacetate (897 μ l, 6.14 mmol). The mixture was refluxed for 24 h. The mixture was cooled to room temperature and then rotary evaporated to dryness. To the residue was added water (40 ml) and the mixture was extracted with chloroform (2 x 60 ml). The chloroform solution was dried over anhydrous sodium sulfate and then concentrated in vacuo. The residue was purified by column chromatography to give the product as a yellow solid (350 mg, 35%). ^1H NMR (CDCl_3): 11.21 (s, 1H), 9.04 (d, $J = 5.1$ Hz, 2H), 7.58 (t, $J = 5.1$ Hz, 1H), 6.98 (s, 1H).

[0312] c. 4-Chloro-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine: A solution of 2-(2-pyrimidinyl)-6-trifluoromethylpyrimidin-4-ol (135 mg, 0.557 mmol) in phosphorus oxychloride (3 ml) was refluxed for 3 h. The mixture was cooled to room temperature and then rotary evaporated to leave a brown oil. The oil was extracted with ethyl acetate (75 ml), and the extracts were washed with water (2x20 ml), and dried over anhydrous sodium sulfate. The ethyl acetate solution was rotary evaporated to dryness to give a yellow solid (120 mg, 83%). ^1H NMR (CDCl_3): 9.08 (d, $J = 4.5$ Hz, 2H), 7.83 (s, 1H), 7.51 (t, $J = 4.5$ Hz, 1H).

[0313] d. 4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride: A mixture of 4-chloro-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine (50 mg, 0.192 mmol) and 2-chloro-5-methoxyaniline hydrochloride (56 mg, 0.288 mmol) in water : ethanol (2:1, 10 ml) was refluxed for 48 h. The mixture was cooled to room temperature and basified with aqueous 2N NaOH to pH 10-12. The resulting precipitate was collected by filtration and purified by column chromatography to give a yellow solid (21 mg, 28%). ^1H NMR (CDCl_3): 9.01 (d, $J = 4.8$ Hz, 2H), 7.71 (s, 1H), 7.67 (s, 1H), 7.44 (t, $J = 4.8$ Hz, 1H), 7.36 (d, $J = 9.3$ Hz, 1H), 7.09 (s, 1H), 6.75 (dd, $J = 3.0, 9.0$ Hz, 1H), 3.86 (s, 3H).

EXAMPLE 215

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(t-butyl)pyrimidine

[0314] a. 4-Chloro-2-(3-pyridinyl)-6-(t-butyl)pyrimidine: The title compound was prepared from 6-tert-butyl-2-pyridin-3-yl-pyrimidin-4-ol (400 mg, 1.75 mmol) and phosphorus oxychloride (10 ml) similar to Example 214c and isolated as a tan solid (333 mg, 77%). ¹H NMR (CDCl₃): 9.68 (d, J = 1.5 Hz, 1H), 8.80-8.76 (m, 1H), 8.73 (dd, J = 1.5, 4.8 Hz, 1H), 7.49-7.44 (m, 1H), 7.28 (s, 1H), 1.41 (s, 9H).

[0315] b. 4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(t-butyl)pyrimidine: The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(tert-butyl)pyrimidine (50 mg, 0.202 mmol) and 2-chloro-5-methoxyaniline hydrochloride (59 mg, 0.303 mmol) similar to Example 117 and isolated as a yellow oil (12 mg, 15%). ¹H NMR (CDCl₃): 9.70 (s, 1H), 8.75-8.68 (m, 2H), 8.13 (d, J = 2.7 Hz, 1H), 7.42-7.39 (m, 1H), 7.31 (d, J = 8.7 Hz, 1H), 7.11 (s, 1H), 6.65 (s, 1H), 6.60 (dd, J = 3.0, 8.7 Hz, 1H), 3.87 (s, 3H), 1.39 (s, 9H).

EXAMPLE 216

4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine

[0316] To a solution of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.144 mmol) in methanol (20 ml) was added concentrated hydrochloric acid (200 ul). The mixture was hydrogenated over palladium (5 wt% on activated carbon) at 50 psi for 6 h. The mixture was filtered through Celite. The filtrate was rotary evaporated to dryness and purified by column chromatography to give a brown oil (3 mg, 6%). ¹H NMR (CDCl₃): 7.83 (s, 1H), 7.31 (d, J = 7.8 Hz, 1H), 6.90-6.85 (m, 3H), 6.77 (dd, J = 1.8, 7.8 Hz, 1H), 3.90 (dd, J = 3.0, 10.5 Hz, 1H), 3.81 (s, 3H), 3.29 (d, J = 11.4 Hz, 1H), 2.88-2.80 (m, 2H), 2.25 (d, J = 10.5 Hz, 1H), 1.92 (s, 1H), 1.69-1.56 (m, 4H).

- 136 -

EXAMPLE 217

4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine

[0317] To a stirring solution of 4-(2-chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.131 mmol) in dichloromethane (5 ml) was added m-chloroperbenzoic acid (23 mg, 0.131 mmol). The mixture was stirred at room temperature for 30 minutes and then rotary evaporated to dryness. The residue was purified by column chromatography to give a white solid (19 mg, 36%). ¹H NMR (CDCl₃): 8.35 (d, J = 7.5 Hz, 2H), 8.28 (d, J = 7.5 Hz, 2H), 7.64 (d, J = 2.4 Hz, 1H), 7.40 (d, J = 9.0 Hz, 1H), 7.30 (s, 1H), 6.94 (s, 1H), 6.76 (dd, J = 3.0, 9.0 Hz, 1H), 3.85 (s, 3H).

EXAMPLE 218

4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0318] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 4-chloro-2-methoxy-5-methylaniline (50 mg, 0.289 mmol) similar to Example 117 and isolated as a tan solid (26 mg, 34%). ¹H NMR (CDCl₃): 9.65-9.63 (m, 1H), 8.75-8.70 (m, 2H), 8.18 (s, 1H), 7.46-7.41 (m, 1H), 7.36 (s, 1H), 6.97 (s, 1H), 6.87 (s, 1H), 3.91 (s, 3H), 2.42 (s, 3H).

EXAMPLE 219

4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0319] a. 2,4,5-Trimethylphenylamine: A solution of 5-nitropseudocumene (400 mg, 2.42 mmol) in methanol (30 ml) was hydrogenated over palladium (5 wt% on activated carbon) at 50 psi for 3 h. The mixture was filtered through Celite. The filtrate was rotary evaporated to

- 137 -

leave a light purple residue (246 mg, 75%). ¹H NMR (CDCl₃): 6.82 (s, 1H), 6.50 (s, 1H), 3.42 (s, 2H), 2.16 (s, 3H), 2.14 (s, 3H), 2.11 (s, 3H).

[0320] b. 4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine: The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) 2,4,5-trimethylphenylamine (39 mg, 0.289 mmol) similar to Example 117 and isolated as a yellow solid (21 mg, 30%). ¹H NMR (CDCl₃): 9.70 (d, J = 1.8 Hz, 1H), 8.73-8.70 (m, 1H), 8.67 (dd, J = 1.5, 4.8 Hz, 1H), 7.65 (s, 1H), 7.42-7.38 (m, 1H), 7.11 (s, 1H), 7.10 (s, 1H), 6.51 (s, 1H), 2.29 (s, 3H), 2.27 (s, 3H), 2.22 (s, 3H).

EXAMPLE 220

4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

[0321] The title compound was prepared from a mixture of 4-chloro-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine (50 mg, 0.193 mmol) and 5-amino-2,4-dichlorobenzenol (51 mg, 0.289 mmol) similar to Example 117 and isolated as a white solid (24 mg, 31%). ¹H NMR (DMSO-d₆): 10.80 (s, 1H), 9.90 (s, 1H), 9.39-9.38 (m, 1H), 8.75-8.72 (m, 1H), 8.57-8.53 (m, 1H), 7.63 (s, 1H), 7.61 (s, 1H), 7.59-7.55 (m, 1H), 7.31 (s, 1H).

EXAMPLE 221

Identification of 4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine and Analogs as Caspase Cascade Activators and Inducers of Apoptosis in Solid Tumor Cells

[0322] Human breast cancer cell lines T-47D and ZR-75-1 were grown according to media component mixtures designated by American Type Culture Collection + 10% FCS (Invitrogen Corporation, Life Technologies division), in a 5% CO₂ -95% humidity incubator at 37 °C. T-47D and ZR-75-1 cells were maintained at a cell density between 50 and 80% confluency at a

- 138 -

cell density of 0.1 to 0.6×10^6 cells/ml. Cells were harvested at 600xg and resuspended at 0.65×10^6 cells/ml into appropriate media + 10% FCS. An aliquot of 45 μ l of cells was added to a well of a 96-well microtiter plate containing 2.5 μ l of a 10% DMSO in RPMI-1640 media solution containing 0.16 to 100 μ M of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine or other test compound (0.016 to 10 μ M final). An aliquot of 22.5 μ l of cells was added to a well of a 384-well microtiter plate containing 2.5 μ l of a 10% DMSO in RPMI-1640 media solution without test compound as the control sample. The samples were mixed by agitation and then incubated at 37 °C for 48 h in a 5% CO₂-95% humidity incubator. After incubation, the samples were removed from the incubator and 25 μ l of a solution containing 14 μ M of *N*-(Ac-DEVD)-*N'*-ethoxycarbonyl-R110 fluorogenic substrate (Cytovia, Inc.; WO99/18856), 20% sucrose (Sigma), 20 mM DTT (Sigma), 200 mM NaCl (Sigma), 40 mM Na PIPES buffer pH 7.2 (Sigma), and 500 μ g/ml lysolecithin (Calbiochem) was added. The samples were mixed by agitation and incubated at room temperature. Using a fluorescent plate reader (Model SpectraMax Gemini, Molecular Devices), an initial reading ($T = 0$) was made approximately 1- 2 min after addition of the substrate solution, employing excitation at 485 nm and emission at 530 nm, to determine the background fluorescence of the control sample. After the 3 h incubation, the samples were read for fluorescence as above ($T = 3$ h).








[0323] Calculation:

[0324] The Relative Fluorescence Unit values (RFU) were used to calculate the sample readings as follows:









[0325]
$$\text{RFU}_{(T=3h)} - \text{Control RFU}_{(T=0)} = \text{Net RFU}_{(T=3h)}$$

[0326] The activity of caspase cascade activation was determined by the ratio of the net RFU value for 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine or other test compound to that of control samples. The EC₅₀ (nM) was determined by a sigmoidal dose-response calculation (Prism 2.0, GraphPad Software Inc.). The caspase activity (Ratio) and potency (EC₅₀) are summarized in Table I:









TABLE I. CASPASE ACTIVITY AND POTENCY

Compd #	Structure	T47 Ratio 48 hr	T47 EC50 (nM)	ZR Ratio 48 hr	ZR EC50 (nM)
A		29.3	433	3.6	140
19		16.7	471	8.9	214
41		7.5	521	1.6	>5000
74		16.4	364	13.1	326
73		1.6	>5000	2.6	10000
131		1.2	>5000	1.2	>5000
164		1.3	>5000	1.3	>5000









- 140 -

125		4.4	3467	8.1	2239
163		1.1	>5000	1.6	>5000
75		16.3	2292	7.8	3276
81		12.7	5052	5.0	5497
44		1.6	>5000	1.9	>5000
45		21.6	329	17.3	387
16		19.5	288	3.3	292
109		3.9	1495	3.6	2594









- 141 -

110		8.9	576	11.6	648
111		4.0	1421	3.9	2569
112		3.3	1860	2.9	2371
113		4.4	1426	4.1	2733
114		3.6	2539	3.7	2800
15		2.5	>5000	2.7	>5000
18		1.9	>5000	7.8	4901
46		1.5	>5000	1.4	>5000









- 142 -

20		1.6	>5000	1.1	>5000
160		1.0	>5000	1.4	>5000
104		1.1	>5000	1.7	>5000
200		1.2	>5000	1.4	>5000
69		14.2	382	10.9	127
92		16.8	363	12.4	285
156		1.7	>5000	4.1	2261
96		1.4	>5000	4.9	2759









- 143 -

93		2.0	>5000	8.2	2032
97		14.5	659	6.7	717
60		1.8	>5000	1.4	>5000
62		1.6	>5000	1.3	>5000
65		1.7	>5000	1.7	>5000
61		1.8	>5000	1.2	>5000
59		1.6	>5000	1.1	>5000
58		1.5	>5000	1.1	>5000

- 144 -

66		1.2	>5000	0.9	>5000
67		1.5	>5000	1.2	>5000
63		8.9	18	9.6	43
38		8.8	19	12.4	69
42		5.8	43	2.5	145
76		8.5	<150	ND ^a	ND
102		8.0	512	7.4	701
85		1.7	>5000	6.7	2834









- 145 -

51		11	29	5.9	56
99		11.1	34	12.0	50
55		8.4	21	7.1	18
80		8.5	64	5.9	243
180		15.1	637	8.7	698
50		10	172	5.3	322
95		11.6	113	8.6	148
82		15.3	254	ND	ND

- 146 -

54		10.5	99	5.9	101
127		10.3	279	5.3	550
188		3.6	429	9.3	271
117		11.3	138	6.3	276
118		8.8	34	9.9	31
194		11.7	107	ND	ND
106		9.0	183	2.4	982
107		8.0	63	5.3	123









- 147 -

21		15.5	<156	12.0	<156
53		20.4	115	6.6	599
52		20.9	108	7.2	98
190		19.3	29	16.7	32
198		3.3	419	1.8	>5000
162		11.3	272	7.3	360
87		19.2	<75	16.9	<75
88		9.8	2315	8.0	1216









- 148 -

89		8.7	2737	9.8	1603
126		12.0	222	5.9	353
123		21.2	763	15.8	725
116		13.7	71	8.8	73
196		11.3	527	ND	ND
68		20.9	247	17.9	130
115		14.2	112	11.6	59
91		11.7	1442	8.8	788









- 149 -

120		8.2	2694	4.5	2397
183		17.3	82	11.5	75
161		10.7	79	5.1	86
122		11.1	44	10.1	192
195		8.4	20	ND	ND
121		11.2	480	9.9	380
128		1.4	>5000	1.5	>5000
124		19.9	720	16.3	688








- 150 -

176		4.5	149	1.9	>5000
179		10.8	1474	1.3	>5000
108		0.9	>5000	2.4	4367
22		1.9	>5000	3.5	6049
130		11.3	295	9.7	167
134		16.5	807	1.5	>5000
133		16.8	1286	4.5	354
157		6.6	43	6.4	197

- 151 -

201		15.7	333	4.8	173
166		8.0	870	3.3	532
206		6.9	802	2.9	583
158		9.5	509	6.0	133
203		10.7	843	3.4	2627
212		5.8	3114	6.1	2781
213		14.1	2685	9.9	1768
147		13.3	335	8.4	204

- 152 -

148		13.8	412	8.4	243
149		16.0	75	10.0	46
150		11.0	135	9.7	133
151		12.0	388	8.6	276
152		14.6	35	8.6	212
153		15.7	43	9.3	146
154		6.1	22	7.9	14

^a. ND, not determined.

- 153 -

Thus, 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (Compound A) and related compounds are identified as potent caspase cascade activators and inducer of apoptosis in solid tumor cells.

EXAMPLE 222

Identification of 4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine as Antineoplastic Compound that Inhibits Cell Proliferation (GI₅₀)

[0327] T-47D cells were grown and harvested as in Example 221. An aliquot of 90 μ l of cells (2×10^4 cells/ml) was added to a well of a 96-well microtiter plate containing 10 μ l of a 10% DMSO in RPMI-1640 media solution containing 1 nM to 100 μ M of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (0.1 nM to 10 μ M final). An aliquot of 90 μ l of cells was added to a well of a 96-well microtiter plate containing 10 μ l of a 10% DMSO in RPMI-1640 media solution without compound as the control sample for maximal cell proliferation (A_{Max}). The samples were mixed by agitation and then incubated at 37°C for 48 h in a 5% CO₂-95% humidity incubator. After incubation, the samples were removed from the incubator and 20 μ l of CellTiter 96 AQUEOUS One Solution Cell Proliferation™ reagent (Promega) was added. The samples were mixed by agitation and incubated at 37 °C for 2-4 h in a 5% CO₂-95% humidity incubator. Using an absorbance plate reader (Model 1420 Wallac Instruments), an initial reading (T=0) was made approximately 1-2 min after addition of the solution, employing absorbance at 490 nm. This determines the possible background absorbance of the test compounds. No absorbance for 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine was found at 490 nm. After the 2-4 h incubation, the samples were read for absorbance as above (A_{Test}).

[0328] Baseline for GI₅₀ (dose for 50% inhibition of cell proliferation) of initial cell numbers were determined by adding an aliquot of 90 μ l of cells or 90 μ l of media, respectively, to wells of a 96-well microtiter plate containing 10 μ l of a 10% DMSO in RPMI-1640 media solution. The samples were

- 154 -

mixed by agitation and then incubated at 37°C for 0.5 h in a 5% CO₂-95% humidity incubator. After incubation, the samples were removed from the incubator and 20 µl of CellTiter 96 AQUEOUS One Solution Cell ProliferationTM reagent (Promega) was added. The samples were mixed by agitation and incubated at 37°C for 2-4 h in a 5% CO₂-95% humidity incubator. Absorbance was read as above, (A_{Start}) defining absorbance for initial cell number used as baseline in GI₅₀ determinations.

[0329] Calculation:

[0330] GI₅₀ (dose for 50% inhibition of cell proliferation) is the concentration where $[(A_{\text{Test}} - A_{\text{Start}}) / (A_{\text{Max}} - A_{\text{Start}})] = 0.5$

[0331] The GI₅₀ (nM) are summarized in Table II

Table II. GI₅₀ in T-47D Cancer Cells

Cell lines	Compound A GI ₅₀ (nM)
T-47D	30

Thus, 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine (Compound A) is identified as antineoplastic compound that inhibits cell proliferation.

EXAMPLE 223

Identification of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine as Antineoplastic Compound that Selectively Inhibits the Proliferation of Breast Cancer Cells(GI₅₀)

[0332] T-47D, ZR-75-1, MX-1, SK-Br-3, MCF-7, MDA-MB-435 (all breast cancer cell lines), Panc-1 (pancreatic cancer cell line), MES-SA (sarcoma cell line), and PC-3 (prostate cancer cell line) cells were grown according to the conditions recommended by American Type Culture Collection. SW620 (colorectal cancer cell line), and P388 (mouse leukemia cell line) were grown according to the conditions provided by the National Cancer Institute. The cell

- 155 -

proliferation assay and the calculations of GI_{50} s were performed as in Example 222 and the results were summarized in Tables III and IV.

Table III. GI_{50} in human breast cancer cell lines.

Cell Line	T47D	ZR 75-1	MCF-7	MDA-MB-435	SK-Br-3
GI_{50} (nM)	30	100	100	4500	60

Table IV. GI_{50} in non-breast cancer cell lines.

Cell Line	PC-3	Panc-1	SW620	P388	MES-SA
GI_{50} (nM)	>10,000	>10,000	>10,000	>10,000	>10,000

Thus, 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine was identified as antineoplastic compound that selectively inhibits the growth of human breast cancer cells.

EXAMPLE 224

Treatment with 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine Leads to Cell Cycle Arrest and Apoptosis in T-47D Cells

[0333] T-47D, a breast cancer cell line, was maintained and harvested as described in Example 221. 5×10^5 Cells were treated with 0.2 μ M of 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine for 48 h at 37 °C. As a control, cells were also incubated with equivalent amount of solvent (DMSO). Cells were harvested at 1,200 rpm and then transferred to 12x75 mm polystyrene tubes. Cells were then resuspended in 500 μ l of 1% Na Citrate, 0.1% Triton X-100, and 50 μ g/ml of propidium iodide and incubated at room temperature for 30 min followed by flow cytometer analysis. All flow cytometry analyses were performed on FACScalibur

- 156 -

(Becton Dickinson) using Cell Quest analysis software. The x-axis plotted the amount of fluorescence and the y-axis is plotted the number of cells with the indicated fluorescence. The T-47D control cell population profile is seen in Fig. 1A and the increase in G2/M DNA content cells that is seen when treated with 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine at 0.2 μ M is seen in Fig. 1B. An increase in the sub-diploid DNA content of cells (Fig 1) is also seen to increase from 2% to 25% with compound treatment. The sub-diploid amount of DNA is indicative of apoptotic cells which have undergone DNA degradation or fragmentation.

EXAMPLE 225

4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine Inhibit the Clonogenic Survival of T47D and MX-1, Solid Tumor Cell Lines

[0334] T47D and MX-1 cells were grown according to the conditions according conditions recommended by American Type Culture Collection. In a well of a 96 well plate, 30,000 cells were seeded and treated with tested compound at the indicated concentrations for 48 hr in a 5% CO₂- 95% humidity incubator at 37°C. Control wells were treated with the same amount of solvent (DMSO) as the compound samples. After the 48 h treatment, the supernatant was removed to a sterile culture tube and the wells washed with phosphate buffered saline, and the adherent cells trypsinized for 5 min. The trypsinized cells were added to the culture supernatant, cells were collected (1,200 rpm, 10 min), washed with phosphate buffered saline, and resuspended in fresh media. The cells were counted for trypan blue negative cells, and the cells diluted with fresh media to 1,000 cells/ml. To a well of a 24-well plate, 0.1 ml of the cell suspension was added along with 1 ml of fresh media (cell suspensions were passed through a 22G needle several times just before plating to form single cell suspensions). Plates are incubated in a 5% CO₂- 95% humidity incubator at 37°C for 7-10 days. Colonies are counted when the sizes reached greater than 50 cells per colony. Cells are washed with phosphate buffered saline, fixed with 100% methanol for 15 min, and then

- 157 -

stained with 0.5% gentian violet for 15 min. Colonies are rinsed with water and the colonies counted and the fraction surviving expressed as the percentage of the number of control colonies.


[0335] The results showed that a 48 hr treatment with 4-(3-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine inhibited the ability of T47D and MX-1 cells to proliferate and their colony forming ability with an IC_{50} of about 100 and 300 nM, respectively.

[0336] Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

- 158 -

WHAT IS CLAIMED IS:

1. A method of treating a disorder responsive to the induction of apoptosis in an animal suffering therefrom, comprising administering to an animal in need of such treatment an effective amount of a compound of Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar₁ and Ar₂ are independently optionally substituted aryl or optionally substituted heteroaryl;

A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, optionally substituted alkyl or optionally substituted cycloalkyl.

2. The method of claim 1, provided that when said disorder is cancer then:

Ar₁ is pyran-2-yl, piperidin-2-yl, piperazin-2-yl, pyrrolidin-2-yl, imidazolidin-2-yl, imidazol-2-yl, indol-3-yl, isoindol-1-yl, quinuclidin-2-yl, morpholin-2-yl, isochroman-2-yl, chroman-2-yl, pyrazolidin-2-yl, pyrazol-2-yl, tetrahydropyran-2-yl, tetrahydro-2H-pyran-2-yl, benzo[b]thien-2-yl, naphtho[2,3-b]thien-2-yl, thianthren-2-yl, isobenzofuran-2-yl, chromen-2-yl, xanthen-2-yl, phenoxanthin-2-yl, 2H-pyrro-2-yl,

- 159 -


pyrrolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalziny, naphthyridinyl, quinoxaliny, cinnoliny, pteridinyl, carbazolyl, β -carboliny, phenanthridinyl, acridinyl, perimidinyl, phenanthroliny, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, phenoxazinyl, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-*a*]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazolyl, 2-oxindolyl, 2-oxobenzimidazolyl, pyridinyl N-oxide, pyrazinyl N-oxide and pyrimidinyl N-oxide; and/or

A is N; and/or

Ar₂ is heteroaryl; and/or

R₃ is cycloalkyl.

3. The method of claim 1, wherein said animal is a mammal.
4. The method of claim 1, wherein A is C-R₂.
5. The method of claim 1, wherein R₃ is hydrogen.
6. The method of claim 1, wherein Ar₂ is optionally substituted and selected from the group consisting of phenyl, naphthyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolyl, isoquinolyl, thienyl, furyl and pyrrolyl.
7. The method of claim 1, wherein said compound has the Formula II:



- 160 -

or a pharmaceutically acceptable salt or prodrug thereof, wherein R_1 - R_3 , and Ar_2 are as defined in claim 1;

B is N or C- R_4 ;

D is N or C- R_5 ;

E is N or C- R_6 ;

F is N or C- R_7 ;

G is N or C- R_8 ; and

R_4 - R_8 are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C_{1-10} alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol;

provided that not more than three of B, D, E, F or G are N.

8. The method of claim 7, wherein R_1 is optionally substituted alkyl, haloalkyl or optionally substituted aryl.

9. The method of claim 7, wherein Ar_2 is optionally substituted phenyl.

10. The method of claim 7, wherein B is N.

11. The method of claim 1, wherein said compound is selected from the group consisting of:

4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Fluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 161 -

4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dichloroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4-Difluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methylanilino)-2-phenyl-6-methylpyrimidine;
4-(4-Methoxyanilino)-2-(2-hydroxyphenyl)-6-methylpyrimidine;
4-(4-(Trifluoromethoxy)anilino)-2-(pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 162 -

4-(4-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-(Trifluoromethoxy)anilino)-2-(2-pyridinyl)-6-*t*-butylpyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-*t*-butylpyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-chloropyrimidine;
4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2-(Trifluoromethyl)benzylamino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

- 163 -

4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 164 -

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 165 -

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;
4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;
4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 166 -

4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine;
4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 167 -

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide;

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

- 168 -

4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine
4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;
6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine;
4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(4-Fluoro-3-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

- 169 -

4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-pyridinyl)pyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(t-butyl)pyrimidine;

4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine; and

4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine.

12. The method of claim 1, wherein said compound is selected from the group consisting of:

- 170 -

4-(3,4,5-Trimethoxyanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3,5-Dichloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(2-Chloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(Indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

- 171 -

4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;


4-(5-Hydroxy-2-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

- 172 -

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
 4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
 4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
 4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
 4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine;
 4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine; and
 4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride.

13. A method for treating or preventing cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar₁ and Ar₂ are independently optionally substituted aryl or optionally substituted heteroaryl;

A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido,

- 173 -

alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, optionally substituted alkyl or optionally substituted cycloalkyl;

provided that wherein said method is for preventing cancer, then

Ar₁ is pyranyl, piperidinyl, piperazinyl, pyrrolidinyl, imidazolidinyl, imidazoliny, indoliny, isoindoliny, quinuclidiny, morpholiny, isochromanyl, chromanyl, pyrazolidiny, pyrazoliny, tetronoyl and tetramoyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthiny, 2*H*-pyrroly, pyrroly, pyraziny, pyrimidiny, pyridaziny, indoliziny, isoindoly, 3*H*-indoly, indoly, indazoly, puriny, 4*H*-quinoliziny, isoquinoly, quinoly, phthalziny, naphthyridiny, quinozaliny, cinnoliny, pteridiny, carbazoly, β-carboliny, phenanthridiny, acridiny, perimidiny, phenanthroliny, phenaziny, isothiazoly, phenothiaziny, isoxazoly, furazany, phenoxaziny, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-*a*]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazoly, 2-oxindoly, 2-oxobenzimidazoly, pyridiny N-oxide, pyraziny N-oxide and pyrimidiny N-oxide; and/or

A is N; and/or

Ar₂ is heteroaryl; and/or

R₃ is cycloalkyl.

14. The method of claim 13, wherein said animal is a mammal.


15. The method of claim 13, wherein said cancer is selected from the group consisting of Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic leukemia, chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, primary

- 174 -

macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, malignant melanoma, choriocarcinoma, mycosis fungoides, head or neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer and prostatic carcinoma.

16. The method of claim 13, wherein said cancer is breast carcinoma.

17. A method for the treatment of drug resistant cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of the Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar₁ and Ar₂ are independently optionally substituted aryl or optionally substituted heteroaryl;

A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido,

- 175 -

alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, optionally substituted alkyl or optionally substituted cycloalkyl.

18. The method of claim 17, wherein said animal is a mammal.

19. The method of claim 13 or 17, additionally comprising administering at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

20. The method of claim 19, wherein said known cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguanzone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

21. The method of claim 13 or 17, additionally comprising treating said animal with radiation-therapy.

22. The method of claim 13 or 17, wherein said compound is administered after surgical treatment of said animal for cancer.

23. The method of claim 1, wherein said disorder is an autoimmune disease.

- 176 -

24. The method of claim 1, wherein said disorder is rheumatoid arthritis.
25. The method of claim 1, wherein said disorder is inflammation.
26. The method of claim 1, wherein said disorder is inflammatory bowel disease.
27. The method of claim 1, wherein said disorder is Crohn's disease.
28. The method of claim 1, wherein said disorder is ulcerative colitis.
29. The method of claim 1, wherein said disorder is a skin disease.
30. The method of claim 29, wherein said disorder is psoriasis.
31. A compound selected from the group consisting of:
 - 4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(4-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(4-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
 - 4-(3-Benzoyloxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Ethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Cyanoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Acetophenonanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
 - 4-(3-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;

- 177 -

4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Acetoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Nitroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-(Trifluoromethoxy)anilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-(Methylthio)anilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(3-Methoxy-phenoxy)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-5-methoxy-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;
6-Morpholino-4-(3-methoxyanilino)-2-phenyl-pyrimidine;
6-Morpholino-4-(2,5-dimethoxyanilino)-2-phenyl-4-pyrimidine;
4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

- 178 -

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-[3-(4-Bromo-1-methyl-1H-pyrazol-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-(2-Methyl-pyrimidin-4-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Phenylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-(3-Nitrophenyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-(2,3,4,5,6-Pentafluorophenoxy)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 179 -

4-[3-(2-Ethyl-1-phenyl-pyrazolin-5-one-3-yl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-(Phenylsulfone)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-(N-phenylamide)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

- 180 -

4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;

4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 181 -

4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;
4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;
4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Methoxybenzylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Carboxyl-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;

- 182 -

4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;

4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;

4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine;

4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-

(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-piperidino-anilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-

(trifluoromethyl)pyrimidine;

- 183 -

4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine;

4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Chloro-5-methoxyanilino)-2-morpholino-6-(methyl)pyrimidine;

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 184 -

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(2,5-Diethoxy-4-morpholinoanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide;

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-(2,5-Dimethyl-4-hydroxy-anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,5-Dimethyl-4-hydroxy-anilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 185 -

4-(2,5-Dimethyl-4-hydroxy-anilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-aminopyrimidine;

4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine

4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(N-Methyl-3-methoxy-anilino)-2-(2-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(3,5-Dimethoxy-anilino)-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine;

- 186 -

4-(5-Chloro-2-methoxy-anilino)-2-(4-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(3,5-Dimethoxy-anilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(2-Chloro-5-methoxy-anilino)-2-(2,6-dichloro-4-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(2-Methoxy-5-methyl-4-nitro-anilino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(Indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(5-Methoxy-2-methyl-4-nitro-anilino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(3-Trifluoromethyl-1-pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(4,5-Dihydro-2-thiazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(1-Pyrazolyl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(4,5,6,7-Tetrahydro-indazol-1-yl)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

4-(1H-3-Pyrazolyl-amino)-2-(3-pyridinyl)-6-trifluoromethyl-pyrimidine;

6-Methyl-2-(2-pyridinyl)-4-(3-trifluoromethylbenzylamino)-pyrimidine;

6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine;

4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(4-Fluoro-3-methoxy-anilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

- 187 -

4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-pyridinyl)pyrimidine;

4-(5-Chloro-2-methoxy-anilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Methoxy-2-nitro-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Carboxy-2-methoxy-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-nitro-anilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Ethanesulfonyl-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methyl-anilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Ethanesulfonyl-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

- 188 -

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methyl-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-methoxy-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Chloro-2-hydroxy-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-isopropyl-anilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethoxyphenylethylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3,4,5-Trichloroanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(3-Methoxy-dibenzofuran-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(1,5,6-Trimethyl-benzimidazol-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

- 189 -

4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(1,3,4-Triazol-1-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride;
4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(t-butyl)pyrimidine;
4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine; and
4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine.


32. A pharmaceutical composition, comprising the compound of claim 31, and a pharmaceutically acceptable carrier.

- 190 -

33. The pharmaceutical composition of claim 32, further comprising at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

34. The pharmaceutical composition of claim 33, wherein said known cancer chemotherapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

35. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar₁ and Ar₂ are independently optionally substituted aryl or optionally substituted heteroaryl;

A is N or C-R₂;

R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido,


- 191 -

alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and

R₃ is hydrogen, optionally substituted alkyl or optionally substituted cycloalkyl.

Provided that when Ar₂ is aryl, and A is C-R₂, then Ar₁ is, pyranyl, piperidinyl, piperazinyl, pyrrolidinyl, imidazolidinyl, imidazoliny, indolinyl, isoindolinyl, quinuclidinyl, morpholinyl, isochromanyl, chromanyl, pyrazolidinyl, pyrazolinyl, tetronoyl and tetramoyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthinyl, 2*H*-pyrrolyl, pyrrolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalziny, naphthyridinyl, quinoxaliny, cinnoliny, pteridinyl, carbazolyl, β-carboliny, phenanthridinyl, acridinyl, perimidiny, phenanthroliny, phenazinyl, isothiazolyl, phenothiaziny, isoxazolyl, furazanyl, phenoxazinyl, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-a]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazolyl, 2-oxindolyl, 2-oxobenzimidazolyl, pyridinyl N-oxide, pyrazinyl N-oxide and pyrimidinyl N-oxide;

36. The pharmaceutical composition of claim 35, wherein said compound has the Formula II:



or a pharmaceutically acceptable salt or prodrug thereof, wherein R₁-R₃, and Ar₂ are as defined in claim 35;

B is N or C-R₄;

D is N or C-R₅;

E is N or C-R₆;

- 192 -

F is N or C-R₇;

G is N or C-R₈; and

R₄-R₈ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol;

provided that not more than three of B, D, E, F or G are N.

37. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound selected from the group consisting of:

- 4-(3-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(2-Fluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(4-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(3,5-Dichloroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(2,4-Difluoroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(2-Methylanilino)-2-phenyl-6-methylpyrimidine;
- 4-(4-Methoxyanilino)-2-(2-hydroxyphenyl)-6-methylpyrimidine;
- 4-(4-(Trifluoromethoxy)anilino)-2-(pyridinyl)-6-(trifluoromethyl)pyrimidine;
- 4-(3-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 193 -

4-(3-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-(Trifluoromethyl)anilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Fluoroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dichloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Fluoroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-(Trifluoromethyl)anilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Chloroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 194 -

4-(4-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(3-Chloroanilino)-2-(2-pyridinyl)-6-(methoxymethyl)pyrimidine;
4-(4-(Trifluoromethoxy)anilino)-2-(2-pyridinyl)-6-*t*-butylpyrimidine;
4-(4-Methoxyanilino)-2-(2-pyridinyl)-6-*t*-butylpyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-chloropyrimidine;
4-(3-Methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Fluoroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2-(Trifluoromethyl)benzylamino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Methoxyanilino)-2-(3-methylphenyl)-6-(methoxymethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Dimethylaminoanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(3-Isopropylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;
4-(3-Methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 195 -

4-(3-Methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

4-(2,5-Dimethoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(2-methyl-1,3-thiazol-4-yl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-6-(methoxymethyl)-2-(3-methylphenyl)pyrimidine;

4-(2,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

- 196 -

4-(3,4-Methylenedioxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-phenyl-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Methoxy-5-phenoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-[2-Methyl-5-(carboxymethylester)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 197 -

4-(2-Methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Fluoro-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Ethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Ethylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Methoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Chloro-5-methoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Methylmercaptoanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(3-Hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(3-Hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;

4-(3,5-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)-pyrimidine hydrochloride;

- 198 -

4-(2,5-Diethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Carboxyl-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,6-Dimethylanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Methoxy-5-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Cyano-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Chloro-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Methoxy-5-trifluoromethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-[6-(trifluoromethyl)-3-pyridinyl]-6-(trifluoromethyl)pyrimidine;
4-(5-Bromo-2-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

- 199 -

4-(2-Bromo-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-[3-Methyl-5-(trifluoromethyl)anilino]-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Chloro-2-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(4-Chloro-2,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;

4-(2,4-Dichloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Acetyl-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(2-Methyl-5-nitroanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

4-(5-Amino-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;

N-[4-Methyl-3-(2-pyridin-3-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

6-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]nicotinamide;

N-[4-Methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]-4-morpholino-benzamide;

4-Chloro-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

4-Methoxy-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;

- 200 -

4-Chloromethyl-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;
4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(2-pyridin-4-yl-6-(trifluoromethyl)pyrimidin-4-ylamino)-phenyl]benzamide;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,5-Dimethyl-4-hydroxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4-Chloro-2,5-dimethoxyanilino)-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(3-Trifluoromethoxyanilino)-2-(2-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine
4-(4-Chloro-2,5-dimethoxyanilino)-2-(4-pyridinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Hydroxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,5-Dimethoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

- 201 -

4-(5-Chloro-2-methoxyanilino)-2-(4-pyridinyl)-6-trifluoromethylpyrimidine;

6-Methyl-4-(3-phenoxyanilino)-2-(2-pyridinyl)pyrimidine;

4-(3-Chloroanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(4-Fluoro-3-methoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3-Isopropoxyanilino)-6-methyl-2-(2-pyridinyl)pyrimidine;

4-(3,4-Dimethoxyanilino)-6-trifluoromethyl-2-(2-pyridinyl)pyrimidine;

4-(5-Chloro-2-methoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

4-(5-Hydroxy-2-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Chloro-5-hydroxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Methoxy-5-methylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

4-(2-Hydroxy-5-isopropylanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;

- 202 -

4-(3,5-Dimethoxyanilino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Cyano-4,5-dimethoxyanilino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(3-pyridinyl)-6-(*t*-butyl)pyrimidine;
4-(4-Chloro-2-methoxy-5-methylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,4,5-Trimethylanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine; and
4-(2,4-Dichloro-5-hydroxyanilino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine.

38. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound selected from the group consisting of:

4-(3,4,5-Trimethoxyanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3,5-Dichloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(2-Chloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;

- 203 -

4-(3,4,5-Trimethoxyanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3,5-Dichloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(2-Chloroanilino)-2-(*N*-pyrrolyl)-1,3,5-triazine;
4-(3-Methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Methoxy-5-methylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(2,5-Dimethylanilino)-2,6-di(2-pyridinyl)pyrimidine;
4-(Indol-4-ylamino)-2-(3-pyridinyl)-6-(trifluoromethyl)pyrimidine;
4-(Indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2,6-Dimethoxypyridin-3-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Methoxy-pyridin-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(4,6-Dimethoxy-pyrimidin-2-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Hydroxy-5-methyl-pyrimidin-4-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(Indol-4-ylamino)-2-(2-pyridinyl)-6-trifluoromethylpyrimidine;
4-(2-Methyl-indol-5-ylamino)-2-(3-pyridinyl)-6-trifluoromethylpyrimidine;
4-(3-Methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;

- 204 -

4-(3,4-Dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(5-Methoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(2-Chloro-5-methoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3,4-Methylenedioxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine;
4-(3-Hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2,5-Dimethyl-4-hydroxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4-Chloro-2,5-dimethoxyanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(4,5-Dimethoxy-2-methylanilino)-2-(2-pyrazinyl)-6-(trifluoromethyl)pyrimidine hydrochloride;
4-(2-Methoxy-5-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Methoxy-2-nitroanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(3,5-Dimethoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Carboxy-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(2-Methoxy-5-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Hydroxy-2-methylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(2-Chloro-5-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;

- 205 -

4-(2,5-Dimethylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-methoxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Chloro-2-hydroxyanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(5-Hydroxy-2-isopropylanilino)-2-(2-pyrazinyl)-6-trifluoromethylpyrimidine;
4-(2-Chloro-5-methoxy-anilino)-2-(2-pyrimidinyl)-6-trifluoromethylpyrimidine hydrochloride;
4-(3-Methoxyanilino)-2-(2-piperidinyl)-6-(trifluoromethyl)pyrimidine; and
4-(2-Chloro-5-methoxyanilino)-2-(4-pyridinyl-N-oxide)-6-(trifluoromethyl)pyrimidine.

39. The pharmaceutical composition of any one of claims 35-38, further comprising at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.


40. The pharmaceutical composition of claim 39, wherein said known cancer chemotherapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

- 206 -



SUBSTITUTED 2-ARYL-4-ARYLAMINOPYRIMIDINES AND
ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF
APOPTOSIS AND THE USE THEREOF

ABSTRACT OF THE DISCLOSURE

The present invention is directed to substituted 2-aryl-4-arylamino-pyrimidine and analogs thereof, represented by the general Formula I:



wherein A, Ar₁, Ar₂, R₁ and R₃ are defined herein. The present invention also relates to the discovery that compounds having Formula I are activators of caspases and inducers of apoptosis. The compounds of this invention may be used to induce cell death in a variety of clinical conditions in which uncontrolled growth and spread of abnormal cells occurs.

Figure 1A**Figure 1B**




Figure 2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/47498

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/50, 31/53, 31/505

US CL : 514/241, 245, 246, 247, 256

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/241, 245, 246, 247, 256

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95/06473 A2 (NIPPON KAYAKU KABUSHIKI KAISHA) 09 March 1995, see the entire document.	1-40
X	WO 98/38171 A1 (SIGNAL PHARMACEUTICALS, INC.) 03 September 1998, see the entire document.	1-40

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 MAY 2002

Date of mailing of the international search report

30 MAY 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3930

Authorized Officer

KEVIN E. WEDDINGTON

Telephone No. (703) 308-0196

stk65



Request: post66m-282 from stk65@srv06

Options:

Title: /tmp/typemail.11351 ##### ####

Fri Jul 11 10:37:35 EDT 2003

From ychen@cas.org Fri Jul 11 09:11:51 2003
Received: from srv01.cas.org (srv01 [134.243.50.9])
by srv06.cas.org (8.11.6+Sun/m4_8.10.2/CAS_CLIENT-1.18) with ESMTP id h6BDBp41153
for <book@srv06.cas.org>; Fri, 11 Jul 2003 09:11:51 -0400 (EDT)
Received: from yyc56mwu.cas.org (yyc56mwu [134.243.140.152])
by srv01.cas.org (8.12.9/8.12.9/CAS_MAIL_HUB-3.02) with ESMTP id h6BDBnRS024524;
Fri, 11 Jul 2003 09:11:50 -0400 (EDT)
Received: (from yyc56@localhost)
by yyc56mwu.cas.org (8.11.6+Sun/m4_8.11.6/CAS_CLIENT-1.18) id h6BDBn517939;
Fri, 11 Jul 2003 09:11:49 -0400 (EDT)
Date: Fri, 11 Jul 2003 09:11:49 -0400 (EDT)
From: yyc56 <ychen@cas.org>
Message-Id: <10307110911.AA17937@cas.org>
Subject: Borrowing Journals or Patents
To: book@cas.org
Content-Length: 1034

Fri Jul 11 09:07:45 EDT 2003

JOURNAL OR PATENT REQUEST

<u>NAME</u>	<u>EXT</u>	<u>DEPT</u>	<u>ROOM</u>	<u>LOGON ID</u>	<u>PRIM SYS</u>
CHEN, YEN Y.	3593	56	4323A	yyc56	srv25

PUBLICATION TITLE: Handbook of Polymer Testing: Physical Methods
[In: Plast. Eng. (N. Y.), 1999]
ISBN 0-8247-0171-2
edited by Roger Brown

YEAR: 1999

VOLUME:

ISSUE:

PAGES:

CODEN (if available):

OTHER (e.g. Patent type or number, title, author of article):

☐ Please mail the piece to my office.
☒ I will be down to pick up the piece.

If you have a preferred format check the one you need:

☐ Assigned Copy
☒ Original
☐ Photocopy

>>> If this item is not immediately available, we will contact you. <<<<

Return to Stacks by _____

From jowens@cas.org Fri Jul 11 09:45:13 2003
Received: from srv01.cas.org (srv01 [134.243.50.9])
by srv06.cas.org (8.11.6+Sun/m4_8.10.2/CAS_CLIENT-1.18) with ESMTTP id h6BDjC40064
for <book@srv06.cas.org>; Fri, 11 Jul 2003 09:45:12 -0400 (EDT)
Received: from jro53mwu.cas.org (jro53mwu [134.243.120.117])
by srv01.cas.org (8.12.9/8.12.9/CAS_MAIL_HUB-3.02) with ESMTTP id h6BDjBRS005089;
Fri, 11 Jul 2003 09:45:11 -0400 (EDT)
Received: (from jro53@localhost)
by jro53mwu.cas.org (8.11.6+Sun/m4_8.11.6/CAS_CLIENT-1.18) id h6BDjBo04261;
Fri, 11 Jul 2003 09:45:11 -0400 (EDT)
Date: Fri, 11 Jul 2003 09:45:11 -0400 (EDT)
From: jro53 <jowens@cas.org>
Message-Id: <10307110945.AA4260@cas.org>
Subject: Borrowing Journals or Patents
To: book@cas.org
Content-Length: 1253

Fri Jul 11 09:43:22 EDT 2003

JOURNAL OR PATENT REQUEST

NAME	EXT	DEPT	ROOM	LOGON ID	PRIM SYS
OWENS, JAMES R.	2319	53	3294A	jro53	srv25
PUBLICATION TITLE: Biochemical and Biophysical Research Communications					
YEAR: 2001					
VOLUME: 284					
ISSUE: 4					
PAGES: 863-869					
CODEN (if available): BBRCA9					
OTHER (e.g. Patent type or number, title, author of article):					

AN 135:193214 CA
TI Transport of Fluorescein in MDCKII-MRP1 Transfected Cells and
mrp1-Knockout Mice
AU Sun, Haiying; Johnson, Dennis R.; Finch, Rick A.; Sartorelli, Alan
C.; Miller, Donald W.; Elmquist, William F.

☐ Please mail the piece to my office.
☒ I will be down to pick up the piece.

If you have a preferred format check the one you need:

☐ Assigned Copy
☐ Original
☒ Photocopy
>>> If this item is not immediately available, we will contact you. <<<<

Return to Stacks by _____
